arXiv:1204.4505v2 [math-ph] 10 May 2012

Standard Modules, Induction and the Structure of the Temperley-Lieb Algebra

DAVID RIDOUT AND YVAN SAINT-AUBIN

(David Ridout) Department of Theoretical Physics
Research School of Physics and Engineering;
and Mathematical Sciences Institute; Australian National University
Canberra, ACT 2600
Australia

david.ridout@anu.edu.au

(Yvan Saint-Aubin) Département de Mathématiques et de Statistique
Université de Montréal
Québec, Canada, H3C 3J7.

saint@dms.umontreal.ca

Abstract

The basic properties of the Temperley-Lieb algebra TL_n with parameter $\beta = q + q^{-1}$, $q \in \mathbb{C} \setminus \{0\}$, are reviewed in a pedagogical way. The link and standard (cell) modules that appear in numerous physical applications are defined and a natural bilinear form on the standard modules is used to characterize their maximal submodules. When this bilinear form has a non-trivial radical, some of the standard modules are reducible and TL_n is non-semisimple. This happens only when q is a root of unity. Use of restriction and induction allows for a finer description of the structure of the standard modules. Finally, a particular central element $F_n \in \mathsf{TL}_n$ is studied; its action is shown to be non-diagonalisable on certain indecomposable modules and this leads to a proof that the radicals of the standard modules are irreducible. Moreover, the space of homomorphisms between standard modules is completely determined. The principal indecomposable modules are then computed concretely in terms of standard modules and their inductions. Examples are provided throughout and the delicate case $\beta = 0$, that plays an important role in physical models, is studied systematically.

1 Introduction

The Temperley-Lieb algebras are key objects both in mathematics and physics. Temperley and Lieb [1] introduced them as complex associative algebras that arose in their study of transfer matrix approaches to (planar) lattice models. This family of algebras, indexed by a positive integer n and a complex number β, spread quickly through the physics community where it underlies the study of Potts models [2], ice-type models [3], and the Andrews-Baxter-Forrester models [4]. That Temperley-Lieb algebras play a fundamental role in our modern understanding of phase transitions cannot be overstated. These algebras were subsequently rediscovered by Jones [5] who used them to define what is now known as the Jones polynomial in knot theory. The Temperley-Lieb algebras are also intimately connected with the representation theory of the symmetric groups through their realisation as natural quotients of the (type A) Hecke algebras (see [6] for example).

As is usual in physical applications, it is the representation theory of the Temperley-Lieb algebra TL_n which is the main focus of attention. Indeed, Temperley and Lieb's original contribution takes place in a 2^n-dimensional representation commonly used by physicists for the study of spin chains. Such representations still form an active direction of research in mathematical physics. There are, in addition, somewhat smaller representations that are perhaps more natural to consider including, in particular, those which we shall refer to in what follows as *link representations*. From these, one obtains quotients that have come to be described as being *standard*. It is well-known that these standard representations are irreducible for almost all values of the parameter β and that for such β, the (finite-dimensional) representations of TL_n are completely reducible. However, it is a curious fact that the β which are thereby excluded consist, to a large degree, of the parameter values that are of most interest to physicists.

This failure of complete reducibility is also well-known and there have been significant efforts by physicists and mathematicians to understand the representation theory for these exceptional values of β. There is a continuing interest in this quest due to the current reinvigoration of the study of logarithmic conformal field theories. To explain, the lattice models studied by physicists are believed to have limits (as $n \to \infty$ say) in which the model should be replaced by a field theory possessing conformal invariance. The representations upon which the lattice model is founded are supposed to be replaced by representations of the quantised algebra of infinitesimal conformal transformations, the Virasoro algebra (see [7] for a recent attempt to formalise this). Moreover, the corresponding conformal field theory is now believed not to be a minimal model, as the original paradigm suggested, but rather some logarithmic version thereof [8–10]. Here, "logarithmic" indicates that the Virasoro algebra representations involved also fail to be completely reducible. Conjecturing results about these logarithmic theories through the study of their lattice counterparts remains a popular approach.

Perhaps the first to seriously address the structure of Temperley-Lieb representations for all $\beta \in \mathbb{C}$ was Martin. In [2], he devotes two chapters to obtaining an explicit construction of the *principal indecomposable* representations of the Temperley-Lieb algebra. The arguments are rather involved and rely heavily upon intricate combinatorics and a detailed study of a collection of primitive idempotents introduced by Wenzl [11]. Shortly thereafter, and independently, Goodman and Wenzl [12] applied a similarly detailed study of these idempotents to prove explicit results concerning the structure of the *blocks* (two-sided ideals) of the Temperley-Lieb algebras. The proofs require a long series of elementary but tedious lemmata. Unfortunately, they exclude the case when $\beta = 0$ which is of considerable physical interest. Nevertheless, their methods lead to explicit and highly non-trivial descriptions for the radicals of the blocks.

We should also mention the well-known contribution of Westbury [13]. His article approaches the representation theory of the Temperley-Lieb algebras from a more algebraic, and less combinatorial, perspective. First, a sufficient condition for complete reducibility in given whose form is very familiar to conformal field theorists. This criterion is the non-degeneracy of an invariant bilinear form acting on the standard representations and Westbury computes the determinant of this form explicitly (the analogous result for the Virasoro algebra is, of course, the formula of Kac for the determinant of the form defined on Verma modules). The technique employed involves recursion relations and was suggested by older work of James and Murphy [14]. (The recursion Westbury employs contains a mistake, as is illustrated by the example he gives.) The rest of Westbury's article addresses what happens when complete reducibility fails. The method involves using induction and restriction to determine abstractly the spaces of homomorphisms between standard representations (although there is mention of certain explicit constructions along the lines of Martin). Proofs are often minimal, incomplete or, in one case, referred to Martin's book. Crucial points are therefore left without complete arguments: The exactness of the sequences satisfied by the restriction and induction of standard modules, the existence of non-trivial isomorphisms between certain standard modules when q is a root of unity, and the construction and completeness of the set of principal indecomposable modules.

While the above makes our dissatisfaction with Westbury's article evident, his algebraic approach to the Temperley-Lieb algebra provides, in our opinion, an excellent road-map to learning this particular corner of representation theory. Indeed, our motivation for writing this article derives in large part from a desire to have a pedagogical and (mostly) self-contained summary of Temperley-Lieb theory, including detailed proofs. We believe that this will be of significant value to mathematical physicists working on lattice models and conformal field theories, as well as provide novice representation-theorists with an excellent worked example over $\mathbb{C}$ which explicitly illustrates some of the difficulties of non-semisimple associative algebras.

Of course, there are many possible approaches to Temperley-Lieb theory and we certainly do not claim that ours is in any way the best. Indeed, we have tried to minimise the level of sophistication required wherever possible while still introducing the most basic tools that are indispensable for non-semisimple representation theory. To illustrate our prejudices in this regard (and to borrow from John Cardy [15]), we mention that the word 'quiver' has just made its only appearance. Other elegant alternatives include deducing the results from their Hecke algebra analogues (see [6] for example) or using Schur-Weyl duality and quantum group representation theory (see [16] for a recent sketch in this direction). We also mention a powerful category-theoretic approach due to Graham and Lehrer [17] in which structural results for TL_n follow as special cases of their investigations into the *affine* Temperley-Lieb algebras.

The article begins in Section 2 with a review of the diagrammatic and algebraic definitions of the Temperley-Lieb algebras, proving their equivalence using a standard combinatorial argument. Here, we follow the seminal work of Jones [5]. We note, in particular, his result concerning a canonical form for monomials constructed from the standard Temperley-Lieb generators. This turns out to be crucial for the analysis of inducing standard representations in Section 6. We remark that we make a "heretical" choice for Temperley-Lieb diagrams, orienting them at ninety degrees to that customarily found in the literature (see [18] for a precedent illustrating this heresy). This choice facilitates the translation between the diagrammatic and algebraic depictions of multiplication: A diagram drawn on the left of another object corresponds to a left-action on this latter object. We also find it notationally convenient for the following section.

Section 3 then introduces the link representations and their quotients, the standard representations. Various basic, but essential, results are proven for the latter, roughly following the beautiful work of Graham and Lehrer on cellular algebras [19] (the standard representations are cell representations in their formalism). Particular attention is paid to the problematic case $\beta = 0$ in which there is a single standard representation upon which the usual invariant bilinear form vanishes identically.

The hard work begins in Section 4 in which we compute the determinant of an invariant bilinear form on the standard representations. Here, we follow Westbury [13] in using module restriction to block-diagonalise the Gram matrices and deduce recursion relations for the diagonalising matrices. (One can also use idempotents to derive such recursion relations; see [17] for example.) As mentioned above, the recursion relation that Westbury gives is incorrect and we discuss in detail the appropriate refinements that have to be made. The determinant formula then leads to the well-known result concerning the generic semisimplicity of the Temperley-Lieb algebras. Section 5 then uses the proof of the determinant formula to compute the dimension of the kernel of the Gram matrix and thence the dimensions of the radicals and irreducible quotients of the standard representations. By utilising pictures known as Bratteli diagrams, we motivate the conjecture that these radicals are themselves irreducible (when non-trivial).

Sections 6 and 7 are devoted to proving this conjecture. The first details what is obtained upon performing the induced module construction on a standard representation of TL_n to obtain a representation of TL_{n+1}. This result is stated in Westbury, but without proof. We then use these induced representations, in the second, to demonstrate the existence of certain homomorphisms between standard representations. This relies crucially upon a computation which shows that a particular element, which we call F_n, of the centre of TL_n, acts in a non-diagonalisable fashion upon appropriately chosen induced modules. The existence of these homomorphisms is enough to prove the conjecture that the radicals of the standard modules are irreducible (or trivial). To our knowledge, this irreducibility was first proven in [17]; our elementary proof appears to be quite different.

The analysis to this point leaves several questions unanswered, one of which is whether we have determined a complete set of mutually non-isomorphic irreducible representations. Answering questions like these require a little more sophistication and so we turn, in Section 8, to the consideration of the principal indecomposable representations. For this we employ, once again, the induced module technique developed in Section 6 to concretely determine the structure of these representations in terms of that of the standard representations. We remark that the proofs rely heavily on an extremely convenient property of the central element F_n (and that this property is not shared by the central element that appears in Westbury). After a brief summary reporting what has been proven, we conclude with two appendices. The first defines F_n and proves the properties that we require. The second lists a selection of standard representation-theoretic results that are needed. The text is liberally peppered throughout with examples chosen to highlight the theory being developed. We hope that the reader will find them useful for understanding this beautiful corner of representation theory.

We end this introduction with a list of common symbols and terms.

Glossary of Terms and Symbols

$\mathcal{M}_n$ and $\mathcal{M}_{n,p}$: link modules	beginning of Section 3
$\mathcal{V}_{n,p} = \mathcal{M}_{n,p}/\mathcal{M}_{n,p+1}$: standard modules	Equation (3.3)
$\mathcal{R}_{n,p} \subset \mathcal{V}_{n,p}$: radical of $\langle \cdot, \cdot \rangle_{n,p}$ on $\mathcal{V}_{n,p}$	Equation (3.10) and Proposition 3.3
$\mathcal{L}_{n,p} = \mathcal{V}_{n,p}/\mathcal{R}_{n,p}$: irreducible quotient of the standard modules	after Proposition 3.3
$\mathcal{P}_{n,p}$: principal indecomposable module with quotient $\mathcal{L}_{n,p}$	beginning of Section 8
$\mathcal{M}\!\downarrow$: restriction of $\mathcal{M}$ from TL_n to TL_{n-1}	Proposition 4.1
$\mathcal{M}\!\uparrow$: induction of $\mathcal{M}$ from TL_n to TL_{n+1}	beginning of Section 6
basis $\mathcal{S}_{n,p}$ for the induced module $\mathcal{V}_{n,p}\!\uparrow$	before Proposition 6.3
$\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_{n,p}$: bilinear form on $\mathcal{V}_{n,p}$	before Lemma 3.1
(n,p) is critical if $q^{2(n-2p+1)} = 1$	after the proof of Corollary 4.2
critical lines, critical strips of the Bratteli diagram	after Corollary 5.2
$d_{n,p}$: number of (n,p)-link states	before Equation (2.8)
F_n and $f_{n,p}$: central element of TL_n and its eigenvalues	Appendix A and Proposition A.2
$G_{n,p}$: Gram matrices ($\langle \cdot, \cdot \rangle_{n,p}$ in the basis of link states)	beginning of Section 4
Jones' normal form and reverse normal form	Proposition 2.2
simple link in a link state	before Lemma 6.1
admissible link state	before Lemma 6.2
a symmetric pair $V_{n,p}$ and $V_{n,p'}$	beginning of Section 7

2 Diagrams and Presentations

Let us define an n-diagram algebra, for n a positive integer, as follows. First, draw two parallel vertical lines and mark n points on each. The $2n$ points obtained are then connected pairwise by n links (arcs) which do not intersect and which lie entirely between the two vertical borders. This gives what we will call an *n-diagram*. Second, form the complex vector space generated formally by the set of all n-diagrams. Third, endow this vector space with a multiplication given by concatenating two n-diagrams, replacing each interior loop by a factor $\beta \in \mathbb{C}$, and identifying (and then removing) the two interior vertical borders. For example, for $n=3$,



(2.1)

This multiplication defines, for each $\beta \in \mathbb{C}$, an associative algebra called the n-diagram algebra. It is easily checked that this algebra has a unit given by the diagram



We note here for future reference that much of what follows is simplified algebraically upon making the identification $\beta = q + q^{-1}$ with $q \in \mathbb{C}^{\times}$. Of course this implies that the resulting theory must be invariant under the exchange of q with q^{-1}.

The n-diagram algebra is in fact isomorphic to the Temperley-Lieb algebra TL_n, as we shall see. The latter algebra is abstractly defined as being generated by a unit $\mathbf{1}$ and elements u_i, $i = 1,2,\ldots,n-1$, satisfying

$$u_i^2 = \beta u_i, \qquad u_i u_{i\pm 1} u_i = u_i \qquad \text{and} \qquad u_i u_j = u_j u_i \quad \text{if } |i-j| > 1. \tag{2.2}$$

Indeed, if we make the identifications

$\mathbf{1} =$



and $u_i =$



, (2.3)

then we can verify explicitly the properties of the unit (which are clear) and the defining relations



$$u_i^2 = \qquad = \beta \qquad = \beta u_i, \tag{2.4a}$$



$$u_i u_{i-1} u_i = \qquad = \qquad = u_i, \tag{2.4b}$$



and $$u_i u_j = \qquad = \qquad = u_j u_i \qquad (|i-j| > 1). \tag{2.4c}$$

The verification of $u_i u_{i+1} u_i = u_i$ is just (2.4b) viewed upside-down. More formally, note that the Temperley-Lieb algebra TL_n has an automorphism specified by $\mathbf{1} \mapsto \mathbf{1}$ and $u_i \mapsto u_{n-i}$. This corresponds to reflecting our diagrams about a horizontal line. Finally, the map $u_i \mapsto -u_i$ defines an isomorphism between the two algebras TL_n with parameters β and $-\beta$.

This shows that the n-diagram algebra is a quotient of the abstract Temperley-Lieb algebra — the diagram algebra might satisfy further independent relations. To show that there are no further relations, hence that we have the isomorphism of algebras claimed above, we only need to show that the dimensions of the diagram algebra and the Temperley-Lieb algebra coincide. To show this, let us first consider the “half-diagrams” obtained from an n-diagram by cutting vertically down the middle. Each half then has n marked points, but only some of these will still be connected by “links”. If a half-diagram has p links, then there will be $n-2p$ points which are not connected to anything. The latter points will be referred to as *defects* and a half-diagram with n points and p links will be called an (n,p)*-link state*. An example with $n=7$ and $p=3$, hence 1 defect, is given by



We will always orient our link states so that the links and defects face to the right.

An n-diagram can now be cut in half and reassembled into a $(2n,n)$-link state by rotating the incorrectly oriented half so that it lies below the other, and then rejoining the defects of each half as they were joined in the original n-diagram:



As this procedure is obviously reversible, this establishes a bijection between n-diagrams and $(2n,n)$-link states.

The set of all (n,p)-link states is in turn in bijection with the increasing walks on $\mathbb{Z}^2$ from $(0,0)$ to $(n-p,p)$ which avoid crossing the diagonal (m,m). Here, "increasing" means that the walker may only move up or to the right at each step. This bijection is easy to describe: Reading an (n,p)-link state from top to bottom, the walker moves up at the k-th step if the k-th marked point closes a link. Otherwise, the walker moves right. For the example above with $n=7$, $p=3$,



The walk will never cross the diagonal because we cannot close a link without first opening it. Conversely, given such an increasing path, we follow it backwards starting from $(n-p,p)$ and construct the corresponding (n,p)-link state from bottom to top: Every time we move down, we open a link. Every time we move left, we close the newest open link, if one exists to be closed. If one does not, then that marked point becomes a defect. As the walk is never above the diagonal, there are never more down moves than left moves remaining, so we are never left with an open link that cannot be closed. The bijection is now clear.

This proves that the number of (n,p)-link states is equal to the number of increasing walks on $\mathbb{Z}^2$ from $(0,0)$ to $(n-p,p)$ which avoid crossing the diagonal. In particular, the dimension of the n-diagram algebra is therefore the number of increasing walks on $\mathbb{Z}^2$ from $(0,0)$ to (n,n) which avoid crossing the diagonal. We will now establish the same result for the abstract Temperley-Lieb algebra TL_n. Here we follow the seminal paper of Jones [5].

First, define a *word* to be a monomial in the Temperley-Lieb generators, $u_{i_1}u_{i_2}\cdots u_{i_k}$ for example. A word is said to be *reduced* if we cannot use the relations (2.2) to rewrite it with fewer generators. The key insight into converting a given word into reduced form is contained in the following result.

Lemma 2.1. *In any reduced Temperley-Lieb word $u_{i_1}u_{i_2}\cdots u_{i_k}$, the maximal index $m=\max\left\{i_j \ : \ j=1,\ldots,k\right\}$ occurs only once.*

Proof. We use induction on the maximal index m of our reduced word, the case $m=1$ being obvious. Suppose then that we have a reduced word in which u_m appears twice or more, so our word has the form $\cdots u_m U u_m \cdots$, in which the maximal index m' of U is less than m. U must also be reduced (otherwise our word would not be), so we may assume that its maximal index appears only once. Now if $m' < m-1$, then u_m commutes with every generator appearing in U, so we can bring the two u_m's together and use $u_m^2 = \beta u_m$ to reduce our word even further. But, $m' = m-1$ means that u_m commutes with every generator in U except u_{m-1}, so this generator appearing only once in U means that we can sandwich it between the two u_m's and use $u_m u_{m-1} u_m = u_m$. Both possibilities contradict the assumption that our word is reduced, so we conclude that u_m can only appear once. ∎

This simple argument allows one to *order* reduced words by pushing the maximal index as far as possible to the right. More specifically, if U is a reduced word and m the maximal index appearing then we claim that it is possible to commute the u_i so that we have the form

$$U = U' \left(u_m u_{m-1} \cdots u_{m-\ell}\right), \tag{2.5}$$

where U' is a reduced word whose maximal index is less than m. If there were a gap in the sequence of indices following U', then the element after the gap would commute with the generators to the left of the gap (up to u_m), so could be relocated to the left of u_m (and absorbed in U'). If the sequence of indices following U' did not decrease, then one could use $u_i^2 = \beta u_i$ or $u_i u_{i-1} u_i = u_i$ to further reduce the word (contradiction). The claim is therefore established.

Since U' is a shorter reduced word with a smaller maximal index, the same arguments apply, allowing us to write it as U'' times another such uniformly decreasing sequence of generators. Induction then leads us to Jones' normal form for reduced Temperley-Lieb words:

Proposition 2.2 (Jones' Normal Form)**.** *Any reduced Temperley-Lieb word $U \in \mathsf{TL}_n$ may be written in the form*

$$U = (u_{j_1} u_{j_1-1} \cdots u_{k_1})(u_{j_2} u_{j_2-1} \cdots u_{k_2}) \cdots (u_{j_r} u_{j_r-1} \cdots u_{k_r}), \tag{2.6}$$

where $0 < j_1 < \cdots < j_{r-1} < j_r < n$ and $0 < k_1 < \cdots < k_{r-1} < k_r < n$. Similarly, any reduced word may also be written as

$$U = (u_{j_1} u_{j_1+1} \cdots u_{k_1})(u_{j_2} u_{j_2+1} \cdots u_{k_2}) \cdots (u_{j_r} u_{j_r+1} \cdots u_{k_r}), \tag{2.7}$$

with $n > j_1 > j_2 > \cdots > j_r > 0$ and $n > k_1 > k_2 > \cdots > k_r > 0$.

Proof. The increasing nature of the j_i follows from the fact that they are the maximal indices of the subwords $\ldots, U'', U', U$. For the k_i, note that if $k_i \geqslant k_{i+1}$ for some i, then we could commute the u_{k_i} appearing at the end of its decreasing sequence to the right until it bumped up against the $u_{k_i+1} u_{k_i}$ appearing in the next decreasing sequence. The defining relations (2.2) again give us a contradiction to our word being reduced. Thus, we must have $0 < k_1 < \cdots < k_{r-1} < k_r < n$ as well. The proof of the second form is similar. ■

We shall refer to words satisfying (2.6) as elements in Jones' normal form and those satisfying (2.7) as being in reverse Jones' normal form.

The point of this theory is to establish that every monomial in the Temperley-Lieb generators is (up to factors of β) equal to a ordered reduced word of the form (2.6), with strictly increasing j_i and k_i. The dimension of the algebra TL_n is then bounded above by the number of such ordered reduced words.[1] Note that if these ordered reduced words were linearly independent, then this bound would become an equality. Our next task is to show that this number coincides with the number of increasing walks on $\mathbb{Z}^2$ from $(0,0)$ to (n,n) which avoid crossing the diagonal. Then, we will have shown that TL_n has a quotient (the n-diagram algebra) whose dimension is equal to our upper bound on the dimension of TL_n. It then follows that the bound is an equality, so the set of ordered reduced words is in fact a basis.

Naturally, this is achieved by constructing a bijection. This is done by stripping the indices j_i and k_i from the ordered reduced words (2.6) of TL_n and encoding them as a walk on $\mathbb{Z}^2$ as follows:

$$\begin{aligned}&(u_{j_1} u_{j_1-1} \cdots u_{k_1})(u_{j_2} u_{j_2-1} \cdots u_{k_2}) \cdots (u_{j_r} u_{j_r-1} \cdots u_{k_r})\\ \longrightarrow &\Big\langle (0,0) \to (j_1,0) \to (j_1,k_1) \to (j_2,k_1) \to (j_2,k_2) \to \cdots \to (j_r,k_{r-1}) \to (j_r,k_r) \to (n,k_r) \to (n,n) \Big\rangle.\end{aligned}$$

---

[1] When $\beta = 0$, we have the slight modification that every monomial in the Temperley-Lieb generators is either equal to a uniquely ordered reduced word of the form (2.6), with strictly increasing j_i and k_i, or is identically zero. The conclusion remains valid.

(The empty word corresponding to the unit **1** in TL_n is encoded as $\langle(0,0) \to (n,0) \to (n,n)\rangle$.) For example, in TL_9 we have

$$(u_1)\,(u_4u_3u_2)\,(u_6u_5u_4u_3)\,(u_8u_7) \qquad \longrightarrow$$



.

The walks so obtained cannot cross the diagonal because $j_{i+1} > j_i \geqslant k_i$, and they are increasing because $j_i < j_{i+1}$ and $k_i < k_{i+1}$. Conversely, any increasing path from $(0,0)$ to (n,n) is entirely determined by its corners. The coordinates of these corners may be used to construct a reduced word of the form (2.6) and it is easy to check that the j_i and k_i so defined satisfy $j_i < j_{i+1}$ and $k_i < k_{i+1}$. Avoiding crossing the diagonal translates into $j_i \geqslant k_i$.

This bijection therefore completes the proof that the dimension of the n-diagram algebra coincides with that of the abstract Temperley-Lieb algebra TL_n (this dimension is obviously finite). As noted above, this is sufficient to conclude that these algebras are in fact isomorphic. It is not difficult now to complete this circle of ideas and actually compute the dimensions of these algebras by counting the increasing paths from $(0,0)$ to (n,n) which avoid crossing the diagonal.

In fact, we shall generalise slightly and count the increasing paths from $(0,0)$ to $(n-p,p)$ which avoid crossing the diagonal. This more general computation then gives the number of (n,p)-link states. Let this number be $d_{n,p}$. Since an increasing path ending at $(n-p,p)$ must pass through either $(n-p-1,p)$ or $(n-p,p-1)$ (but not both!), we obtain the recursion relation

$$d_{n,p} = d_{n-1,p} + d_{n-1,p-1}. \tag{2.8}$$

With the "boundary values" $d_{n,0} = 1$ (the only increasing path which never goes up is the one which only moves to the right) and $d_{2p-1,p} = 0$ (the paths are not allowed to cross the diagonal), this completely determines the numbers $d_{n,p}$. Its solution is easily verified to be

$$d_{n,p} = \binom{n}{p} - \binom{n}{p-1}, \tag{2.9}$$

where we understand, as usual, that $\binom{n}{-1} = 0$. It follows that the n-diagram algebra and the Temperley-Lieb algebra TL_n have dimension

$$\dim \mathsf{TL}_n = d_{2n,n} = \binom{2n}{n} - \binom{2n}{n-1} = \frac{1}{n+1}\binom{2n}{n}. \tag{2.10}$$

This is of course the n-th Catalan number.

To summarise, we have proven the following result in this section.

Theorem 2.3. *The abstract Temperley-Lieb algebra* TL_n *and the algebra of* n*-diagrams are isomorphic (for given* $n \in \mathbb{Z}_+$ *and* $\beta \in \mathbb{C}$*). The dimensions of these algebras are given by Equation* (2.10)*.*

We mention that combining the result on dimensions with the "cutting" of n-diagrams into two (n, p)-link states leads to the curious identity

$$d_{2n,n} = \sum_{p=0}^{\lfloor n/2 \rfloor} d_{n,p}^2. \tag{2.11}$$

This only requires observing that the two link states must have the same number of links p so that each defect of the first may be joined to a defect of the second.

3 Standard Representations

The (n,p)-link states are not just convenient for combinatorial arguments. They in fact admit a very natural action of the Temperley-Lieb algebra TL_n. More precisely, if we let $\mathcal{M}_n$ denote the complex span of the (n,p)-link states (over all p), then $\mathcal{M}_n$ is naturally a left TL_n-module under the concatenation of diagram with link state. We will refer to this module as the *link module*. An example should serve to make the action clear:



(3.1)

Equation (3.7) provides an example in which a loop is closed upon concatenation. We note that the number of defects need not be conserved under the TL_n-action. However, this action can only close defects in pairs, so the number of defects never increases. Equivalently, the number of links never decreases under the TL_n-action.

We can therefore identify TL_n-submodules $\mathcal{M}_{n,p} \subseteq \mathcal{M}_n$ which are spanned by the (n,p')-link states with $p' \geqslant p$. This gives us an obvious filtration

$$0 \subset \mathcal{M}_{n,\lfloor n/2 \rfloor} \subset \cdots \subset \mathcal{M}_{n,1} \subset \mathcal{M}_{n,0} = \mathcal{M}_n, \tag{3.2}$$

whose consecutive quotients will be denoted by

$$\mathcal{V}_{n,p} = \frac{\mathcal{M}_{n,p}}{\mathcal{M}_{n,p+1}}. \tag{3.3}$$

The $\mathcal{V}_{n,p}$ are therefore TL_n-modules in which the action is by concatenation when the number of links (and defects) is conserved and zero otherwise. We shall refer to them as the *standard modules*. As vector spaces, they are spanned by the (n,p)-link states (we shall often forget to distinguish an element of $\mathcal{M}_{n,p}$ from its coset in $\mathcal{V}_{n,p}$ when it is not crucial), so their dimensions are the $d_{n,p}$ computed in Equation (2.9). For example, $\mathcal{V}_{n,0}$ is always one-dimensional, spanned by the link state with no links (all defects). The Temperley-Lieb generators all act trivially on $\mathcal{V}_{n,0}$ (they would all close two defects), except the unit of course, which acts as the identity.

It is natural to ask at this point if the $\mathcal{V}_{n,p}$ are irreducible as TL_n-modules. We will see, eventually, that the answer is generically "yes". The story is subtle however and will not be completed until the end of Section 4. We will approach this irreducibility in a manner familiar to physicists, by studying the non-degeneracy of an invariant bilinear form. Let us therefore recall that we have oriented our link states so that the links and defects point to the right. This facilitates a left action of the Temperley-Lieb algebra. We could also introduce the reflections of the link states across a vertical line. The links and defects would then point to the left, and we would have a natural right action of TL_n on these reflected link states. This suggests that we might be able to treat reflected link states as linear functionals acting on the link states, hence that combining a reflected link state with a standard link state may lead to an interesting bilinear pairing.

We therefore define, on each $\mathcal{V}_{n,p}$, a form $\langle \cdot,\cdot \rangle \equiv \langle \cdot,\cdot \rangle_{n,p}$ as follows. If x and y are two (n,p)-link states, then $\langle x,y \rangle$ is computed by reflecting x across a vertical line and identifying its vertical border with that of y. The value $\langle x,y \rangle$ is then non-zero only if every defect of x ends up being connected to one of y; when this is so, $\langle x,y \rangle = \beta^m$ where m is the number of closed loops so obtained. This is then extended bilinearly[2] to all of $\mathcal{V}_{n,p}$. As examples, we compute that in $\mathcal{V}_{4,1}$,



Note that this bilinear form is symmetric — $\langle x,y \rangle$ and $\langle y,x \rangle$ are just reflections of one another in the pictorial interpretation.

Consider now the reflection of an n-diagram about a vertical line. This will be another n-diagram, so reflection defines a linear map from TL_n to itself. This obviously preserves the multiplication except that the order will be reversed — this map is an antiautomorphism of TL_n.[3] We will therefore regard it as an adjoint, denoting it by $U \mapsto U^\dagger$. It is clear from Equation (2.3) that $\mathbf{1}^\dagger = \mathbf{1}$ and $u_i^\dagger = u_i$. This definition is natural and useful. Indeed, it shows that the bilinear forms we have defined turn out to be *invariant* with respect to the TL_n-action.

Lemma 3.1. *The bilinear form $\langle \cdot,\cdot \rangle$ on $\mathcal{V}_{n,p}$ satisfies*

$$\langle x,Uy \rangle = \langle U^\dagger x,y \rangle \qquad \text{for all } U \in \mathsf{TL}_n \text{ and } x,y \in \mathcal{V}_{n,p}. \tag{3.5}$$

To see this, we merely note that the two sides of (3.5) are identical when expressed in terms of diagrams and link states.

We introduce a convenient notation $\bigl| x \quad y \bigr|$ for the unique n-diagram which, when cut in half vertically, decomposes into the (n,p)-link state x and the vertical reflection of the (n,p)-link state y. Extending linearly, we obtain a map $\bigl| \cdot \quad \cdot \bigr|$ from $\mathcal{V}_{n,p} \times \mathcal{V}_{n,p}$ into TL_n (in fact we obtain such a map for each p, but we will not bother to distinguish them). The utility of the bilinear form we have defined stems from the following relation.

Lemma 3.2. *If $x,y,z \in \mathcal{V}_{n,p}$, then*

$$\bigl| x \quad y \bigr| z = \langle y,z \rangle x. \tag{3.6}$$

---

[2] What follows can easily be adapted to the case where one extends sesquilinearly. We have chosen bilinearity for simplicity.

[3] We will take this adjoint to be linear as we are considering $\langle \cdot,\cdot \rangle$ as a bilinear form. If one prefers a sesquilinear form, then the adjoint should be taken to be antilinear. Note however, that this conflicts with the defining relations (2.2) unless β is real.

Proof. Observe first that linearity allows us to assume that x, y and z are in fact (n,p)-link states. If any of the defects of y are closed by a link in z, then $\langle y,z\rangle = 0$ by definition, so the right-hand side of (3.6) vanishes. As the defects of x join those of y, this will close two defects of x, leading to an additional link in $\big|x \quad y\big|z$. But then, the resulting link state will vanish in $\mathcal{V}_{n,p}$ due to its definition as a quotient, hence the left-hand side of (3.6) likewise vanishes.

It remains to check the case in which none of the defects of y are closed by a link in z. But then it is clear that the left-hand side of (3.6) will be proportional to x. The proportionality constant is then given by β to the power of the number of loops in the concatenation of $\big|x \quad y\big|$ and z. Since x contributes no loops, this constant is just $\langle y,z\rangle$ as required. ∎

We illustrate this deceptively simple result with an example in $\mathcal{V}_{7,3}$: Compare



$= \beta$ with $\langle \quad , \quad \rangle = \beta.$ (3.7)

We remark that it is tempting (and sometimes useful) to think of $\big|x \quad y\big|$ as the combination $xy^\dagger$, where the "adjoint" of the link state y refers likewise to its vertical reflection. Then, one could "prove" Lemma 3.2 as follows:

$$\big|x \quad y\big|z = \left(xy^\dagger\right)z = x\left(y^\dagger z\right) = \langle y,z\rangle x. \tag{3.8}$$

However, caution should be exercised with such manipulations. If x and y belong to $\mathcal{V}_{n,p'}$ and z is an (n,p)-link state (with $p' \neq p$), then (3.6) and (3.8) do not make sense — the bilinear form is no longer defined. Nevertheless, the product $\big|x \quad y\big|z$ may be non-zero (in $\mathcal{M}_n$) and not even proportional to x. For example,



(3.9)

We will therefore refrain from trying to define an adjoint of a link state (what one means is of course a linear functional in the dual vector space).

We can now understand how this bilinear form will help to decide the irreducibility of the $\mathcal{V}_{n,p}$. For this, we consider the *radical* $\mathcal{R}_{n,p}$ of the bilinear form on $\mathcal{V}_{n,p}$:

$$\mathcal{R}_{n,p} = \left\{x \in \mathcal{V}_{n,p} \ : \ \langle x,y\rangle = 0 \text{ for all } y \in \mathcal{V}_{n,p}\right\}. \tag{3.10}$$

The invariance (Lemma 3.1) of the bilinear form implies that the radical $\mathcal{R}_{n,p}$ is a submodule of $\mathcal{V}_{n,p}$.

Proposition 3.3. *If $\langle \cdot, \cdot \rangle$ is not identically zero on $\mathcal{V}_{n,p}$, then $\mathcal{V}_{n,p}$ is cyclic and indecomposable. Moreover, $\mathcal{V}_{n,p}/\mathcal{R}_{n,p}$ is then irreducible. Equivalently, $\mathcal{R}_{n,p}$ is the unique maximal proper submodule of $\mathcal{V}_{n,p}$.*

(We shall discuss the case when $\langle \cdot, \cdot \rangle$ is identically zero shortly.)

Proof. Since $\langle \cdot, \cdot \rangle$ is not identically zero, there exist $y, z \in \mathcal{V}_{n,p}$ such that $\langle y, z \rangle = 1$. Then, Equation (3.6) says that for every $x \in \mathcal{V}_{n,p}$, we may form $\left| x \quad y \right| \in \mathsf{TL}_n$ to obtain

$$\left| x \quad y \right| z = \langle y, z \rangle x = x. \tag{3.11}$$

Thus, z generates $\mathcal{V}_{n,p}$, proving that this module is cyclic. Now, any $z \notin \mathcal{R}_{n,p}$ has such a partner y, so every such z is a generator of $\mathcal{V}_{n,p}$. It follows that every non-zero element of $\mathcal{V}_{n,p}/\mathcal{R}_{n,p}$ generates this quotient, hence that the quotient is irreducible.

Suppose then that $\mathcal{V}_{n,p}$ is decomposable: $\mathcal{V}_{n,p} = A \oplus B$. Then, $\mathcal{R}_{n,p} = (\mathcal{R}_{n,p} \cap A) \oplus (\mathcal{R}_{n,p} \cap B)$ and it follows from the irreducibility of $\mathcal{V}_{n,p}/\mathcal{R}_{n,p}$ that either A or B must be a submodule of $\mathcal{R}_{n,p}$. Without loss of generality, take $B \subseteq \mathcal{R}_{n,p}$. Then, there exists a non-zero $z \in A$ which is not in $\mathcal{R}_{n,p}$. Such a z cannot generate $\mathcal{V}_{n,p} = A \oplus B$, a contradiction. ■

We will find it convenient to denote the quotient $\mathcal{V}_{n,p}/\mathcal{R}_{n,p}$ by $\mathcal{L}_{n,p}$, even when $\langle \cdot, \cdot \rangle$ vanishes identically. In this latter case, $\mathcal{L}_{n,p}$ is the trivial module $\{0\}$; otherwise, $\mathcal{L}_{n,p}$ is irreducible. For this reason, we shall often refer to the $\mathcal{L}_{n,p}$ as the irreducibles, understanding that one should exclude the exceptional case $\mathcal{L}_{2p,p}$ at $\beta = 0$.

To prove that $\mathcal{V}_{n,p}$ is irreducible, it is therefore enough to show that the radical $\mathcal{R}_{n,p}$ is zero. Equivalently, we must show that the bilinear form we have defined on $\mathcal{V}_{n,p}$ is non-degenerate. This will be the strategy of Section 4. Note however that $\mathcal{R}_{n,p} \neq \{0\}$ only implies that $\mathcal{V}_{n,p}$ is reducible when $\langle \cdot, \cdot \rangle \neq 0$ (that is, when $\mathcal{R}_{n,p} \neq \mathcal{V}_{n,p}$). In case $\langle \cdot, \cdot \rangle = 0$, we cannot decide the irreducibility of $\mathcal{V}_{n,p}$ using Proposition 3.3.

However, when $\beta \neq 0$, $\langle x, x \rangle = \beta^p \neq 0$ for all (n, p)-link states x, so the bilinear form is non-zero. Indeed, if $\beta = 0$ but $n \neq 2p$ (so there is at least one defect), then the form is non-zero because we may choose x and y so as to form a “snake”:

$$\langle x,y\rangle =$$



$$= 1. \tag{3.12}$$

However, when $\beta = 0$ and there are no defects ($n = 2p$), the bilinear form vanishes identically. It is therefore useful to renormalise it, defining

$$\langle x,y\rangle' = \lim_{\beta\to 0} \frac{\langle x,y\rangle}{\beta} \qquad (x,y \in \mathcal{V}_{2p,p}), \tag{3.13}$$

where the forms appearing within the limit are those of the $\mathcal{V}_{2p,p}$ with $\beta \neq 0$. Since $\langle x,y\rangle$ is a polynomial in β with vanishing constant coefficient, $\langle x,y\rangle'$ is defined — in fact, when x and y are $(2p,p)$-link states, it is 1 when a single loop is formed and 0 otherwise. Moreover, $\langle\cdot,\cdot\rangle'$ inherits bilinearity, symmetry and invariance from $\langle\cdot,\cdot\rangle$. Note that this renormalised bilinear form is *not* identically zero because x and y may be chosen so as to obtain

$$\langle x,y\rangle' = \lim_{\beta\to 0} \frac{1}{\beta}$$



$$= \lim_{\beta\to 0} \frac{\beta}{\beta} = 1. \tag{3.14}$$

We cannot immediately apply Proposition 3.3 to the radical of this renormalised form because the proof relied crucially upon Lemma 3.2, special to the form $\langle\cdot,\cdot\rangle$. However, we have the following replacement:

Lemma 3.4. *Let x and y be $(2p,p-1)$-link states, so that they have precisely two defects. Denote by x' and y' the respective $(2p,p)$-link states formed by closing these defects. Then for $\beta = 0$,*

$$\bigl|x \quad y\bigr| z = \langle y',z\rangle' x', \qquad \text{for all } z \in \mathcal{V}_{2p,p}. \tag{3.15}$$

This extends linearly to all $x,y \in \mathcal{V}_{2p,p-1}$.

Proof. Note first that the defects of y must be closed by a link in z (assuming by linearity that the latter is a $(2p,p)$-link state). Thus, $\lvert x \quad y \rvert z$ will be proportional to the link state x' obtained by closing the defects of x. The constant of proportionality is given by β to the power of the number of loops formed by y and z. This is 0 (since $\beta = 0$) unless there are no such loops, in which case it is 1. However, this matches the value of $\langle y', z \rangle'$, as closing the defects of y leads to an additional loop which is dealt with by dividing by β in (3.13). ■

If $\mathcal{R}'_{2p,p}$ denotes the radical of the renormalised $\beta = 0$ form $\langle \cdot, \cdot \rangle'$, we can now mimic the proof of Proposition 3.3 to obtain the analogous result.

Proposition 3.5. *The form $\langle \cdot, \cdot \rangle$ is identically zero if and only if $\beta = 0$ and $n = 2p$. Then, $\mathcal{V}_{2p,p}$ is cyclic, indecomposable and has an irreducible quotient $\mathcal{V}_{2p,p}/\mathcal{R}'_{2p,p}$.*

It remains only to remark that this quotient is never zero because we have shown that $\mathcal{R}'_{2p,p} \neq \mathcal{V}_{2p,p}$ (the form $\langle \cdot, \cdot \rangle'$ is never identically zero).

We conclude this section with a study of whether the $\mathcal{V}_{n,p}$, and their irreducible quotients $\mathcal{L}_{n,p} = \mathcal{V}_{n,p}/\mathcal{R}_{n,p}$ are all mutually distinct as TL_n-modules (up to isomorphism of course). The fact that $\mathcal{V}_{n,p}$ and $\mathcal{V}_{n,p'}$ involve different numbers of links, for $p \neq p'$, can be misleading. For example, $\mathcal{V}_{2,1}$ and $\mathcal{V}_{2,0}$ are isomorphic one-dimensional TL_2-modules, when $\beta = 0$ ($\mathbf{1}$ and u_1 are represented by 1 and 0, respectively, on both). Nevertheless, this behaviour is rather exceptional.

Proposition 3.6. *Let $\mathcal{N}$ and $\mathcal{N}'$ be submodules of $\mathcal{V}_{n,p}$ and $\mathcal{V}_{n,p'}$, respectively, where $p > p'$ and $\langle \cdot, \cdot \rangle_{n,p} \neq 0$. Then, the only module homomorphism $\theta \colon \mathcal{V}_{n,p}/\mathcal{N} \to \mathcal{V}_{n,p'}/\mathcal{N}'$ is the zero homomorphism.*

Proof. Let γ be the canonical homomorphism from $\mathcal{V}_{n,p}$ onto $\mathcal{V}_{n,p}/\mathcal{N}$. Choose $y, z \in \mathcal{V}_{n,p}$ such that $\langle y, z \rangle = 1$. Then, for all $x \in \mathcal{V}_{n,p}$,

$$\lvert x \quad y \rvert \theta(\gamma(z)) = \theta\bigl(\gamma(\lvert x \quad y \rvert z)\bigr) = \theta(\gamma(x)). \tag{3.16}$$

But when $p > p'$, $\lvert x \quad y \rvert \theta(\gamma(z)) = 0$, since $\theta(\gamma(z))$ has p' links but left-multiplying by $\lvert x \quad y \rvert$ leads to at least p links. Thus, $\theta(\gamma(x)) = 0$, so $\theta = 0$ as γ is surjective. ■

Putting $\mathcal{N} = \mathcal{N}' = \{0\}$ or $\mathcal{N} = \mathcal{R}_{n,p}$ and $\mathcal{N}' = \mathcal{R}_{n,p'}$ in Proposition 3.6, and using the basic fact that an isomorphism has an inverse, we obtain:

Corollary 3.7. *When $\langle \cdot, \cdot \rangle_{n,p}$ and $\langle \cdot, \cdot \rangle_{n,p'}$ are non-zero,*

$$\mathcal{V}_{n,p} \cong \mathcal{V}_{n,p'} \quad \Rightarrow \quad p = p' \qquad \text{and} \qquad \mathcal{L}_{n,p} \cong \mathcal{L}_{n,p'} \quad \Rightarrow \quad p = p'.$$

As we have seen, $\langle \cdot, \cdot \rangle_{2p,p}$ vanishes identically when $\beta = 0$, so it follows that $\mathcal{V}_{2p,p}$ ($\mathcal{L}_{2p,p}$) could coincide with one of the other $\mathcal{V}_{2p,p'}$ ($\mathcal{L}_{2p,p'}$) in this case. This is what allows the ($\beta = 0$) isomorphism $\mathcal{V}_{2,1} \cong \mathcal{V}_{2,0}$ which we remarked upon above.

We record a related result for future reference:

Proposition 3.8. *Every module homomorphism $\theta \colon \mathcal{V}_{n,p} \to \mathcal{V}_{n,p}$ is a multiple of the identity.*

Proof. When $\langle \cdot,\cdot \rangle_{n,p} \neq 0$, this follows readily by choosing $y,z \in \mathcal{V}_{n,p}$ such that $\langle y,z \rangle = 1$. Then, for all $x \in \mathcal{V}_{n,p}$,

$$\theta(x) = \theta\left(\big| x \quad y \big| z\right) = \big| x \quad y \big| \theta(z) = \langle y, \theta(z) \rangle x. \tag{3.17}$$

For the remaining case, when $\langle \cdot,\cdot \rangle_{n,p} = 0$, we must have $n = 2p$ and $\beta = 0$. However, the renormalised form $\langle \cdot,\cdot \rangle'$ of Equation (3.13) does not vanish identically, so there exist $y', z \in \mathcal{V}_{2p,p}$ such that $\langle y', z \rangle' = 1$. For every $x' \in \mathcal{V}_{2p,p}$, let $x \in \mathcal{V}_{2p,p-1}$ be obtained by cutting an outermost link of x' (say the one closing at $n = 2p$ for definiteness). Form $y \in \mathcal{V}_{2p,p-1}$ from y' in the same fashion. Then, Lemma 3.4 gives

$$\theta(x') = \theta\left(\big| x \quad y \big| z\right) = \big| x \quad y \big| \theta(z) = \langle y', \theta(z) \rangle' x', \tag{3.18}$$

completing the proof. ■

4 Gram Matrices

We can now turn to a study of the irreducibility of the $\mathcal{V}_{n,p}$, based on the non-degeneracy of the invariant bilinear forms $\langle \cdot,\cdot \rangle_{n,p}$. Recall that each $\mathcal{V}_{n,p}$ has a canonical basis given by the (n,p)-link states. With respect to this basis, the corresponding form is represented by a symmetric $d_{n,p} \times d_{n,p}$ matrix which we shall denote by $G_{n,p}$. Such matrices are called *Gram matrices*. For example,

$$G_{4,0} = \begin{pmatrix} 1 \end{pmatrix}, \qquad G_{4,1} = \begin{pmatrix} \beta & 1 & 0 \\ 1 & \beta & 1 \\ 0 & 1 & \beta \end{pmatrix} \qquad \text{and} \qquad G_{4,2} = \begin{pmatrix} \beta^2 & \beta \\ \beta & \beta^2 \end{pmatrix}, \tag{4.1}$$

when we adopt the (respective) ordered bases





and



As the radical $\mathcal{R}_{n,p}$ is represented by the kernel of the Gram matrix $G_{n,p}$, we see that the irreducibility of the $\mathcal{V}_{n,p}$ is equivalent to $\det(G_{n,p}) \neq 0$.[4] Our aim in this section is to compute $\det(G_{n,p})$ explicitly. The strategy is to use restriction to derive a recursion relation for $\det(G_{n,p})$ when the $\mathcal{R}_{n,p}$ all vanish. This turns out to occur for generic $\beta \in \mathbb{C}$, excluding only a countable set. Continuity therefore takes care of the outstanding cases.

Proposition 4.1. *Consider the inclusion of* TL_{n-1} *in* TL_n *(for fixed* β*) given by sending the unit to the unit and the* u_i *with* $i < n-1$ *to their counterparts in* TL_n*. Denote the corresponding restriction of* $\mathcal{V}_{n,p}$ *to a* TL_{n-1}*-module by* $\mathcal{V}_{n,p}\!\downarrow$*. Then, we have an exact sequence of* TL_{n-1}*-modules,*

$$0 \longrightarrow \mathcal{V}_{n-1,p} \longrightarrow \mathcal{V}_{n,p}\!\downarrow \longrightarrow \mathcal{V}_{n-1,p-1} \longrightarrow 0, \tag{4.2}$$

meaning that $\mathcal{V}_{n-1,p}$ *is a submodule of* $\mathcal{V}_{n,p}\!\downarrow$ *and* $\mathcal{V}_{n,p}\!\downarrow / \mathcal{V}_{n-1,p} \cong \mathcal{V}_{n-1,p-1}$*.*

Proof. The inclusion $\mathcal{V}_{n-1,p} \hookrightarrow \mathcal{V}_{n,p}\!\downarrow$ is defined to extend an $(n-1,p)$-link state to an (n,p)-link state by adding a defect at position n. This is clearly an injective homomorphism of TL_{n-1}-modules as the inclusion of TL_{n-1} in TL_n will preserve the defect at n. The quotient $\mathcal{V}_{n,p}\!\downarrow / \mathcal{V}_{n-1,p}$ is then a TL_{n-1}-module with a basis of cosets which are represented by the (n,p)-link states in which n is part of a link.

There is an obvious vector space isomorphism Ψ from $\mathcal{V}_{n,p}\!\downarrow / \mathcal{V}_{n-1,p}$ to $\mathcal{V}_{n-1,p-1}$ obtained by cutting the link which closes at n and then removing the newly-created defect at n. We wish to show that Ψ is in fact an isomorphism of TL_{n-1}-modules, thereby completing the proof. So for a given basis element z of $\mathcal{V}_{n,p}\!\downarrow / \mathcal{V}_{n-1,p}$, let m denote the opening point of the link which closes at n. Applying any u_i $(i < n-1)$ to z usually then gives another such basis element in which n is linked to some m'. It is easy to see that $\Psi(u_i z) = u_i \Psi(z)$ in this case. The only exception occurs if $i = m-1$ and $m-1$ is a defect in z. Then, applying u_{m-1} to z leads to n becoming a defect:

---

[4] We shall defer addressing the exceptional case of $\mathcal{V}_{2p,p}$ with $\beta = 0$ until the end of the section.



(4.3)

In this case, $\Psi(u_{m-1}z) = \Psi(0) = 0$. But, $\Psi(z)$ will have defects at both $m-1$ and m, so applying u_{m-1} will close the defect leading to an extra link. Thus, $u_{m-1}\Psi(z) = 0$ too and Ψ is a homomorphism. ■

Corollary 4.2. *When* $q^{2(n-2p+1)} \neq 1$ *(recall that we write* $\beta = q + q^{-1}$*), the exact sequence* (4.2) *splits, so we have*

$$\mathcal{V}_{n,p}\!\downarrow\; \cong \mathcal{V}_{n-1,p} \oplus \mathcal{V}_{n-1,p-1} \qquad \text{(as } \mathsf{TL}_{n-1}\text{-modules).} \tag{4.4}$$

Proof. This is easy to see by using the eigenvalues of the central element F_{n-1} on TL_{n-1} (see Appendix A). Since F_{n-1} is central, left-multiplication by $F_{n-1} - \lambda\mathbf{1}$ is a homomorphism from any module $\mathcal{N}$ to itself, for any $\lambda \in \mathbb{C}$. Thus, the (generalised) eigenspaces of F_{n-1} on $\mathcal{N}$ are submodules. Applying this to $\mathcal{N} = \mathcal{V}_{n,p}\!\downarrow$ and recalling that $\mathcal{V}_{n-1,p}$ and $\mathcal{V}_{n-1,p-1}$ are indecomposable, we see that F_{n-1} has at most two eigenvalues: $f_{n-1,p}$ with eigenspace $\mathcal{V}_{n-1,p}$ and $f_{n-1,p-1}$ with eigenspace (isomorphic to) $\mathcal{V}_{n-1,p-1}$. If these eigenvalues are distinct, both eigenspaces are submodules and (4.2) splits. It remains to determine when $f_{n-1,p} = f_{n-1,p-1}$. But, Proposition A.2 gives

$$f_{n-1,p-1} - f_{n-1,p} = \left(q - q^{-1}\right)\left(q^{n-2p+1} - q^{-(n-2p+1)}\right). \tag{4.5}$$

If either factor is zero, then $q^{2(n-2p+1)}$ will be 1, contradicting the hypothesis. ■

We introduce a useful definition: Say that the pair (n,p) is *critical* for a given $q \in \mathbb{C}^{\times}$ if $q^{2(n-2p+1)} = 1$. Similarly, any quantity indexed by n and p will be said to be critical if (n,p) is critical. A rephrasing of Corollary 4.2 is therefore that a non-critical restricted module $\mathcal{V}_{n,p}\!\downarrow$ always splits as $\mathcal{V}_{n-1,p} \oplus \mathcal{V}_{n-1,p-1}$.

Suppose now that $\mathcal{V}_{n,p}\!\downarrow$ is non-critical so that there exists a splitting $\psi\colon \mathcal{V}_{n,p}\!\downarrow\; \longrightarrow \mathcal{V}_{n-1,p} \oplus \mathcal{V}_{n-1,p-1}$. Then, if we order the canonical basis of link states of $\mathcal{V}_{n,p}$ so that those of $\mathcal{V}_{n-1,p}$ (which have n as a defect) come first, ψ may be chosen so that it is represented by a matrix of the form

$$U_{n,p} = \begin{pmatrix} \mathrm{id} & V_{n,p} \\ 0 & \mathrm{id} \end{pmatrix}. \tag{4.6}$$

Here, the submatrix $V_{n,p}$ encodes the non-trivial part of the splitting (the embedding of $\mathcal{V}_{n-1,p-1}$ in $\mathcal{V}_{n,p}\!\downarrow$).

At this point, we make an inductive assumption: We suppose that when q is not a root of unity (so (4.2) always splits), the $\mathcal{R}_{n',p'}$ *vanish* for all $n' < n$ and all p' — this is certainly true for $n' \leqslant 2$. With this assumption, the $\mathcal{V}_{n',p'}$ are all irreducible and mutually distinct by Proposition 3.3 and Corollary 3.7.

Lemma 4.3. *If the splitting* ψ *exists, we may define a bilinear form on* $\mathcal{V}_{n-1,p} \oplus \mathcal{V}_{n-1,p-1}$ *by*

$$\langle\!\langle x+x', y+y' \rangle\!\rangle = \left\langle \psi^{-1}\left(x+x'\right), \psi^{-1}\left(y+y'\right) \right\rangle_{n,p}, \qquad \text{for } x,y \in \mathcal{V}_{n-1,p} \text{ and } x',y' \in \mathcal{V}_{n-1,p-1}. \tag{4.7}$$

This form is symmetric and invariant, hence

$$\langle\!\langle x+x', y+y' \rangle\!\rangle = \langle x,y \rangle_{n-1,p} + \alpha_{n,p} \langle x',y' \rangle_{n-1,p-1}, \tag{4.8}$$

for some $\alpha_{n,p} \in \mathbb{C}$.

Proof. This is a well-known argument: A bilinear form induces a map from a module V to its dual by $v \mapsto \langle v, \cdot \rangle$. The invariance of the form translates into this induced map being an intertwiner. When V is irreducible, so is its dual, so Schur's lemma tells us that the induced maps form a one-dimensional vector space, hence so too do the invariant bilinear forms. In the application at hand, V is the direct sum of two non-isomorphic irreducible modules, so there is a two-dimensional space of bilinear forms. Comparing the form on $\mathcal{V}_{n-1,p}$ with that on the direct sum fixes one of the latter's degrees of freedom to be unity. ■

In matrix form, this becomes

$$\begin{aligned} G_{n-1,p} \oplus \alpha_{n,p} G_{n-1,p-1} &= \left(U_{n,p}^{-1}\right)^T G_{n,p} U_{n,p}^{-1} \\ \Rightarrow \qquad G_{n,p} &= U_{n,p}^T \begin{pmatrix} G_{n-1,p} & 0 \\ 0 & \alpha_{n,p} G_{n-1,p-1} \end{pmatrix} U_{n,p}. \end{aligned} \tag{4.9}$$

This is the recurrence relation which we shall use to compute $\det(G_{n,p})$. For this, it is useful to introduce the familiar notation $[m]_q$ for the q-number

$$[m]_q = \frac{q^m - q^{-m}}{q - q^{-1}}, \tag{4.10}$$

with the limiting case for $q = \pm 1$ being $[m]_q = mq^{m-1}$. Note that (n,p) is critical if and only if $[n-2p+1]_q = 0$ or $q = \pm 1$.

Proposition 4.4. *When* $[n-2p+1]_q \neq 0$ *and* $p > 0$, $\alpha_{n,p}$ *is finite and is given by*

$$\alpha_{n,p} = \frac{[n-2p+2]_q}{[n-2p+1]_q}. \tag{4.11}$$

Proof. We will prove this first under the assumption that q is not a root of unity. The general case then follows from continuity.

We begin by writing the Gram matrix in block form. We again choose to order the canonical basis of link states so that those with n a defect come first. Thus,

$$G_{n,p} = \begin{pmatrix} G_{n,p}^{1,1} & G_{n,p}^{1,2} \\ G_{n,p}^{2,1} & G_{n,p}^{2,2} \end{pmatrix}, \tag{4.12}$$

where $G_{n,p}^{1,1}$ is in fact just $G_{n-1,p}$ (removing the defect at n has no effect on the values taken by the bilinear form). Substituting Equations (4.6) and (4.12) into Equation (4.9) now gives two independent (non-trivial) constraints:

$$G_{n,p}^{1,2} = G_{n-1,p} V_{n,p} \tag{4.13a}$$

$$\text{and} \qquad G_{n,p}^{2,2} = (V_{n,p})^T G_{n-1,p} V_{n,p} + \alpha_{n,p} G_{n-1,p-1} \tag{4.13b}$$

$$= (V_{n,p})^T G_{n,p}^{1,2} + \alpha_{n,p} G_{n-1,p-1}, \tag{4.13c}$$

where the last equality uses (4.13a).

Just as we chose the basis of $\mathcal{V}_{n,p}$ so that the link states with n a defect came first, we now refine it so that when n is a defect, the link states with $n-1$ a defect come before those with $n-1$ part of a link. Similarly, when n is part of a link, put those link states with $n-1$ linked to n before the rest. Pictorially, the order is:



This results in $G_{n,p}^{1,2}$, $G_{n,p}^{2,2}$ and $V_{n,p}$ having the (refined) block forms

$$G_{n,p}^{1,2} = \begin{pmatrix} 0 & * \\ G_{n-2,p-1} & * \end{pmatrix}, \quad G_{n,p}^{2,2} = \begin{pmatrix} \beta G_{n-2,p-1} & * \\ * & * \end{pmatrix} \quad \text{and} \quad V_{n,p} = \begin{pmatrix} V_{n,p}^{1,1} & * \\ V_{n,p}^{2,1} & * \end{pmatrix}. \tag{4.14}$$

To explain, the top-left block of $G_{n,p}^{1,2}$ corresponds to the scalar product of link states of types 1 and 3 (in the above order), which clearly gives zero. Similarly, the bottom-left block of $G_{n,p}^{1,2}$ corresponds to the scalar product of link states of types 2 and 3, and it is easy to see that the value will not change if we remove $n-1$ and n from both (cutting one link in each). Finally, the top-left block of $G_{n,p}^{2,2}$ represents the scalar product of two type 3 link states, hence the factor of β.

We substitute the block forms (4.14) and (4.12) (with $n \to n-1$) into Equation (4.13a). The first column then yields two equations:

$$0 = G_{n-2,p} V_{n,p}^{1,1} + G_{n-1,p}^{1,2} V_{n,p}^{2,1} \tag{4.15a}$$

$$\text{and} \qquad G_{n-2,p-1} = G_{n-1,p}^{2,1} V_{n,p}^{1,1} + G_{n-1,p}^{2,2} V_{n,p}^{2,1}. \tag{4.15b}$$

Using Equation (4.13a) (again with $n \to n-1$) on (4.15a) gives

$$V_{n,p}^{1,1} + V_{n-1,p} V_{n,p}^{2,1} = 0, \tag{4.16}$$

since $G_{n-2,p}$ is invertible by assumption ($\mathcal{R}_{n-2,p} = \{0\}$). To simplify Equation (4.15b), we note that $G^{2,1}_{n-1,p}$ is the transpose of $G^{1,2}_{n-1,p}$ and apply (4.12), (4.13b) (both with $n \to n-1$) and then (4.16) to get

$$V^{2,1}_{n,p} = \frac{1}{\alpha_{n-1,p}} \operatorname{id}. \tag{4.17}$$

Here we have used the invertibility of $G_{n-2,p-1}$.

Finally, we look at the top-left block of Equation (4.13c). Substituting the block forms (4.14), applying (4.17) and the invertibility of $G_{n-2,p-1}$ once more, we arrive at a recursion relation for the $\alpha_{n,p}$:

$$\alpha_{n,p} = \beta - \frac{1}{\alpha_{n-1,p}}. \tag{4.18}$$

This simple relation allows us to reduce the computation of any $\alpha_{n,p}$ to that with n smaller. In particular, if $n = 2p$, we have $G_{n,p} = \beta G_{n-1,p-1}$, since $\mathcal{V}_{n-1,p}$ is not defined and $(n-1,p-1)$-link states are lifted to $\mathcal{V}_{n,p}$ by adding a link. This gives the starting point for the recursion, $\alpha_{2p,p} = \beta$. It is easy to check that Equation (4.11) is the unique solution. ■

Since $\det(U_{n,p}) = 1$, Equation (4.9) now yields an explicit recursion relation for the determinants of the Gram matrices:

$$\det(G_{n,p}) = \alpha_{n,p}^{d_{n-1,p-1}} \det(G_{n-1,p}) \det(G_{n-1,p-1}). \tag{4.19a}$$

Here, $d_{n-1,p-1}$ is the dimension of $\mathcal{V}_{n-1,p-1}$ as given in Equation (2.9). The starting points for this recursion are the results

$$\det(G_{n,0}) = 1 \qquad \text{and} \qquad \det(G_{2p-1,p}) = 1. \tag{4.19b}$$

The first follows because $\mathcal{V}_{n,0}$ is spanned by a single link state consisting entirely of defects. The second follows from putting $G_{2p,p} = \beta G_{2p-1,p-1}$ into (4.19a).

It is clear from Equations (4.11) and (4.19) that the determinant of $G_{n,p}$ can only vanish when q is a root of unity. This gives the following fundamental result:

Proposition 4.5. *When q is not a root of unity, the determinant of the Gram matrices of the $\mathcal{V}_{n,p}$ are all non-zero, hence the $\mathcal{V}_{n,p}$ are irreducible TL_n-modules.*

Corollary 4.6. *When q is not a root of unity, TL_n is a semisimple algebra and the $\mathcal{V}_{n,p}$, $0 \leqslant p \leqslant \lfloor n/2 \rfloor$, form a complete set of non-isomorphic irreducible modules.*

Proof. The $\mathcal{V}_{n,p}$ are irreducible by Proposition 4.5, distinct by Corollary 3.7, and form a complete set (meaning that there are no other non-isomorphic irreducibles) because the sum of the squares of their dimensions gives that of TL_n (Equation (2.11)). The result now follows from Proposition B.2. ■

We remark that the irreducibility of the $\mathcal{V}_{n,p}$ for all p is not sufficient to conclude that TL_n is semisimple. A counterexample is TL_2 with $\beta = 0$ for which the left regular representation is reducible but indecomposable, despite both $\mathcal{V}_{2,1}$ and $\mathcal{V}_{2,0}$ being irreducible (they both have dimension 1). The issue here is that, as we saw in Section 3, these two standard modules are in fact isomorphic. The sum of the squares of the dimensions of the *distinct* irreducibles is thus $1 < \dim \mathsf{TL}_2 = 2$, so we cannot deduce semisimplicity. For this, it is sufficient to have both the irreducibility and the inequivalence of the $\mathcal{V}_{n,p}$.

Let us return now to the recursion relation (4.19) for the $\det(G_{n,p})$. Its solution gives us the main result of this section.

Theorem 4.7. *For all n, p and all $\beta = q + q^{-1} \in \mathbb{C}$, the Gram matrix of $\mathcal{V}_{n,p}$ has determinant*

$$\det(G_{n,p}) = \prod_{j=1}^{p} \left(\frac{[n-2p+1+j]_q}{[j]_q} \right)^{d_{n,p-j}}. \tag{4.20}$$

Proof. As in the proof of Proposition 4.4, we may first assume that q is not a root of unity so that the recursion (4.19) is well-defined for all n and p (q general then follows from continuity). Checking Equation (4.19b) is easy, and (4.19a) follows from considering the numerators and denominators separately and applying Equation (2.8) several times. ■

Note that Equation (4.20) is manifestly invariant under $q \leftrightarrow q^{-1}$ because q-numbers are unchanged under this transformation. We remark that it is not entirely obvious from this formula that $\det(G_{n,p})$ is actually finite for all $q \neq 0$, though this is clear from the definition (it is polynomial in β). Here is the most useful consequence of the determinant formula (4.20):

Corollary 4.8. *If (n, p) is critical, then $\mathcal{R}_{n,p} = \{0\}$, so $\mathcal{V}_{n,p}$ is irreducible.*

Proof. Recall that (n, p) critical means that $q^{2(n-2p+1)} = 1$. We let ℓ be the smallest positive integer for which $q^{2\ell} = 1$, so that $n - 2p + 1 = m\ell$ for some positive integer m, and

$$\det(G_{n,p}) = \prod_{j=1}^{p} \left(\frac{[m\ell + j]_q}{[j]_q} \right)^{d_{n,p-j}}. \tag{4.21}$$

The numerator only vanishes when $j = m'\ell$ for some positive integer m' (this also requires $\ell > 1$), but then the denominator vanishes likewise. Now note that

$$[m\ell]_q = \frac{q^{\ell} - q^{-\ell}}{q - q^{-1}} \frac{q^{m\ell} - q^{-m\ell}}{q^{\ell} - q^{-\ell}} = [\ell]_q [m]_{q^{\ell}}, \tag{4.22}$$

so the zero of $[m\ell]_q$ is always first order (because $q^{\ell} = \pm 1$, so $[m]_{q^{\ell}} \neq 0$ for $m \neq 0$). Thus, the zeroes in the numerator of (4.21) precisely cancel those in the denominator, hence $\det(G_{n,p}) \neq 0$. ■

This corollary implies the semisimplicity of TL_n when $\beta = \pm 2$ ($q = \pm 1$) because all (n, p) are then critical. We similarly obtain the semisimplicity of the TL_n with n odd at $\beta = 0$ ($q = \pm \mathrm{i}$). Indeed,

$\ell = 2$ divides $n-2p+1$ in this case, so the (n,p) with n odd are all critical. The inequivalence of the $\mathcal{V}_{n,p}$ follows from Corollary 3.7 as n odd means, in particular, that $n \neq 2p$ for any p. Even though the (n,p) are not critical when $\beta = 0$ and n is even, the tools developed up to now allow for the proof that $\mathcal{V}_{2p,p}$, which then coincides with $\mathcal{R}_{2p,p}$, is irreducible.

Proposition 4.9. *When $\beta = 0$, the radical $\mathcal{R}_{2p,p}$ is irreducible.*

Proof. Since $\mathcal{R}_{2p,p} = \mathcal{V}_{2p,p}$ when $\beta = 0$, we must show that $\mathcal{R}'_{2p,p} = \{0\}$ (Proposition 3.5). Let $G'_{2p,p}$ denote the Gram matrix of the renormalised form $\langle \cdot,\cdot \rangle'$ defined in Equation (3.13). As noted before Proposition 4.5, $G_{2p,p} = \beta G_{2p-1,p-1}$ and $G'_{2p,p} = \lim_{\beta\to 0} G_{2p,p}/\beta = G_{2p-1,p-1}|_{\beta=0}$. For $\beta = 0$, the values of q are $\pm\mathfrak{i}$ and $[m]_q$ is 0 when m is even and ± 1 when it is odd. Therefore,

$$\det\left(G'_{2p,p}\right) = \det\left(G_{2p-1,p-1}\right)\big|_{\beta=0} = \prod_{j=1}^{p-1} \left(\frac{[j+2]_q}{[j]_q}\right)^{d_{2p-1,p-1-j}} \Bigg|_{q=\pm\mathfrak{i}} . \tag{4.23}$$

Each factor of this product with j odd is a power of ± 1. When j is even, we use $[2m]_q = [2]_q [m]_{q^2}$ to see that the factor's numerator and denominator are both polynomials in β with a simple root at $\beta = 0$. Their quotient is therefore non-zero at $\beta = 0$. It follows that $\det\left(G'_{2p,p}\right) \neq 0$, whence the result. ■

5 Explorations at Roots of Unity

The results of Section 4 not only tell us when the radical $\mathcal{R}_{n,p}$ is trivial, but they also tell us its dimension when it is not. Recall that Equation (2.8) amounts to a simple recursion relation for the dimensions of the standard modules:

$$\dim \mathcal{V}_{n,p} = \dim \mathcal{V}_{n-1,p} + \dim \mathcal{V}_{n-1,p-1}. \tag{5.1}$$

The analogue for the $\mathcal{R}_{n,p}$ is given as follows. Let q be a root of unity and let ℓ be the minimal positive integer satisfying $q^{2\ell} = 1$. For given (n,p) then, set

$$n - 2p + 1 = k(n,p)\,\ell + r(n,p), \tag{5.2}$$

where $k(n,p) \in \mathbb{N}$ and $r(n,p) \in \{1, \ldots, \ell-1, \ell\}$. The criticality of (n,p) is therefore equivalent to $r(n,p) = \ell$ when q is a root of unity.

Proposition 5.1. *The dimensions of the radicals $\mathcal{R}_{n,p}$ satisfy the recursion relation*

$$\dim \mathcal{R}_{n,p} = \begin{cases} 0 & \text{if } r(n,p) = \ell, \\ \dim \mathcal{R}_{n-1,p} + \dim \mathcal{V}_{n-1,p-1} & \text{if } r(n,p) = \ell - 1, \\ \dim \mathcal{R}_{n-1,p} + \dim \mathcal{R}_{n-1,p-1} & \text{otherwise,} \end{cases} \tag{5.3}$$

with initial conditions $\dim \mathcal{R}_{n,0} = 0$ *and* $\dim \mathcal{R}_{2p-1,p} = 0$.

Proof. The initial conditions are clear as $G_{n,0} = \begin{pmatrix} 1 \end{pmatrix}$ for all n and $\mathcal{V}_{2p-1,p}$ is not defined for all p. The recurrence when $r(n,p) = \ell$, which is when (n,p) is critical, also follows directly from Corollary 4.8. We may therefore assume that $r(n,p) \neq \ell$. Then, $q^{2(n-2p+1)} = q^{2r(n,p)} \neq 1$ by minimality of ℓ, so Corollary 4.2 applies. We can therefore make a change of bases to bring $G_{n,p}$ to the block diagonal form of Equation (4.9):

$$G_{n,p} = U_{n,p}^T \begin{pmatrix} G_{n-1,p} & 0 \\ 0 & \alpha_{n,p} G_{n-1,p-1} \end{pmatrix} U_{n,p}. \tag{5.4}$$

We want the dimension of the kernel of $G_{n,p}$. But, as $U_{n,p}$ is invertible,

$$G_{n,p} v = 0 \qquad \Longleftrightarrow \qquad \begin{pmatrix} G_{n-1,p} w_1 \\ \alpha_{n,p} G_{n-1,p-1} w_2 \end{pmatrix} = 0, \qquad \text{where } \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = U_{n,p} v. \tag{5.5}$$

Thus when $\alpha_{n,p} \neq 0$, $\ker(G_{n,p}) = \ker(G_{n-1,p}) \oplus \ker(G_{n-1,p-1})$ as vector spaces, whereas when $\alpha_{n,p} = 0$, $\ker(G_{n,p}) = \ker(G_{n-1,p}) \oplus \mathcal{V}_{n-1,p-1}$ as vector spaces. The result now follows from Proposition 4.4. ∎

Corollary 5.2. *The dimensions of the irreducible quotients $\mathcal{L}_{n,p} = \mathcal{V}_{n,p}/\mathcal{R}_{n,p}$ satisfy the recursion relation*

$$\dim \mathcal{L}_{n,p} = \begin{cases} \dim \mathcal{V}_{n,p} & \text{if } r(n,p) = \ell, \\ \dim \mathcal{L}_{n-1,p} & \text{if } r(n,p) = \ell - 1, \\ \dim \mathcal{L}_{n-1,p} + \dim \mathcal{L}_{n-1,p-1} & \text{otherwise,} \end{cases} \tag{5.6}$$

with initial conditions $\dim \mathcal{L}_{n,0} = 1$ *and* $\dim \mathcal{L}_{2p-1,p} = 0$.

$$
\begin{array}{ccccccccc}
 & | & & & & & & & \\
 & (1,0) & & & & & & & \\
(2,1) & | & (2,0) & | & & & & & \\
 & (3,1) & & (3,0) & & & & & \\
(4,2) & | & (4,1) & | & (4,0) & | & & & \\
 & (5,2) & & (5,1) & & (5,0) & & & \\
(6,3) & | & (6,2) & | & (6,1) & | & (6,0) & | & \\
 & (7,3) & & (7,2) & & (7,1) & & (7,0) & \\
(8,4) & | & (8,3) & | & (8,2) & | & (8,1) & | & (8,0) \\
 & \vdots & & \vdots & & \vdots & & \vdots &
\end{array}
$$

Figure 1. An example of the arrangement of pairs called a Bratteli diagram. The critical lines in this picture correspond to $\beta = 0$, hence $\ell = 2$.

We can use Proposition 5.1 and Corollary 5.2 to build up tables of dimensions of radicals and irreducibles. It proves convenient to arrange these tables so that n increases as we go down and p increases in a south-westerly direction (staying constant to the south-east). The important quantity $n-2p+1$ is therefore constant along vertical lines and increasing from left to right. This corresponds to the standard arrangement of the Bratteli diagram of the family of Temperley-Lieb algebras. To illustrate this, we present the first few rows of this diagram in Figure 1. It is useful to mark the critical (n,p) — these form vertical lines on the diagram which we will refer to as *critical lines*. The regions bounded by consecutive critical lines will be referred to as *critical strips*.

When we replace the pairs in the Bratteli diagram by some quantity indexed by n and p, we shall also refer to the result as a Bratteli diagram. In Figure 2, we show the first few rows of the Bratteli diagrams obtained by replacing (n,p) by $\dim \mathcal{R}_{n,p}$ and $\dim \mathcal{L}_{n,p}$ when $\beta = \pm\sqrt{2}$ ($q = \pm e^{i\pi/4}, \pm e^{3\pi i/4}$). The features of these tables are similar for other roots of unity. Notice that the radicals vanish on the critical lines in accordance with Corollary 4.8.

The feature which is of most interest to us here is that the dimensions of the non-critical $\mathcal{R}_{n,p}$ appear to coincide with those of the $\mathcal{L}_{n,p'}$, where p' is obtained from p by reflecting about the critical line immediately to the right. This suggests that when the radical $\mathcal{R}_{n,p}$ does not vanish, it is in fact *irreducible*. Establishing this irreducibility is somewhat difficult and will be the focus of Section 7. For now, we content ourselves with demonstrating this observed coincidence of dimensions.

Proposition 5.3. *Let q be a root of unity and (n,p) be non-critical. Then, $\dim \mathcal{R}_{n,p}$ is equal to $\dim \mathcal{L}_{n,p+r(n,p)-\ell}$ if $p+r(n,p)-\ell \geqslant 0$ and 0 otherwise.*

We remark that the indices (n,p) of $\mathcal{R}$ and $(n,p+r(n,p)-\ell)$ of $\mathcal{L}$ form a *symmetric pair*, a concept introduced in Section 7.



Figure 2. The Bratteli diagrams for the dimensions of the radicals $\mathcal{R}_{n,p}$ and the irreducibles $\mathcal{L}_{n,p}$ when $\beta = \pm\sqrt{2}$, hence $\ell = 4$.

Proof. This is clear for $n = 1$ as then, $\ell > 2$ and the index $p + r(n,p) - \ell = 2 - \ell$ is negative, so that $\dim \mathcal{R}_{n,p} = 0$ as needed. So assume that the proposition is true for $n-1$ and all p. We have to consider four cases for the pairs (n,p), corresponding to whether $(n-1,p)$ and/or $(n-1,p-1)$ lie on critical lines. The position of the pairs $(n-1,p)$, (n,p) and $(n-1,p-1)$, relative to the critical lines, is pictured below in each case.



Case (1)



Case (2)



Case (3)



Case (4)

1) Both $(n-1,p)$ and $(n-1,p-1)$ are non-critical, so $r(n,p) \notin \{1, \ell-1, \ell\}$. Then,

$$\dim \mathcal{R}_{n-1,p} = \dim \mathcal{L}_{n-1,p+r(n-1,p)-\ell} = \dim \mathcal{L}_{n-1,p-1+r(n,p)-\ell} \tag{5.7a}$$

$$\text{and} \qquad \dim \mathcal{R}_{n-1,p-1} = \dim \mathcal{L}_{n-1,p-1+r(n-1,p-1)-\ell} = \dim \mathcal{L}_{n-1,p+r(n,p)-\ell}. \tag{5.7b}$$

Proposition 5.1 now gives $\dim \mathcal{R}_{n,p}$ as the sum of the left-hand sides. To compute the sum of the right-hand sides, we note that $r(n, p+r(n,p)-\ell) = -r(n,p) \pmod{\ell}$, hence Corollary 5.2 gives $\dim \mathcal{L}_{n,p+r(n,p)-\ell}$ for this sum.

2) Only $(n-1, p-1)$ is critical. Then, $r(n,p) = \ell - 1$ and (5.7a) is valid, but instead of (5.7b), we have

$$\dim \mathcal{V}_{n-1,p-1} = \dim \mathcal{L}_{n-1,p-1} = \dim \mathcal{L}_{n-1,p+r(n,p)-\ell}. \tag{5.7c}$$

Now apply Proposition 5.1 and Corollary 5.2 to the sum of (5.7a) and (5.7c).

3) Only $(n-1, p)$ is critical. Now, $r(n,p) = 1$, (5.7b) is valid, and we use

$$\dim \mathcal{R}_{n-1,p} = 0. \tag{5.7d}$$

Adding (5.7b) and (5.7d) then gives the result.

4) Both $(n-1, p)$ and $(n-1, p-1)$ are critical. Thus, $r(n,p) = 1 = \ell - 1$, that is, $\ell = 2$ (and so $\beta = 0$). The result now follows from adding (5.7c) and (5.7d).

■

We remark that there is one (non-trivial) case in which we already know that the radical is irreducible. This is the content of Proposition 4.9 which asserts that $\mathcal{R}_{2p,p}$ is irreducible when $\beta = 0$. Of course, this does not tell us that $\mathcal{R}_{2p,p}$ is isomorphic to $\mathcal{L}_{2p,p-1}$ (since $\ell = 2$ when $\beta = 0$) as Proposition 5.3 would have us believe.

6 Induced Modules

Recall that in Proposition 4.1, we studied the result of restricting the action on a TL_n-module to the action of the subalgebra TL_{n-1} spanned by the n-diagrams in which the two n-th points are joined. Restriction has a close relative which constructs instead a TL_{n+1}-module from a given TL_n-module (subject to the analogous inclusion of TL_n in TL_{n+1}). This is, of course, the *induced module* construction. We therefore consider the induced modules

$$\mathcal{V}_{n,p}\!\uparrow\; = \mathsf{TL}_{n+1} \otimes_{\mathsf{TL}_n} \mathcal{V}_{n,p}. \tag{6.1}$$

These are TL_{n+1}-modules in which the action is given by left-multiplication: $a\,(b \otimes z) = (ab) \otimes z$ for all $a, b \in \mathsf{TL}_{n+1}$ and $z \in \mathcal{V}_{n,p}$. The subtlety of this definition lies in the subscript on the "$\otimes$" which informs us that the tensor product is "permeable" to elements of TL_n. More precisely, this means that $ab \otimes z = a \otimes bz$ for all $a \in \mathsf{TL}_{n+1}$, $b \in \mathsf{TL}_n$ and $z \in \mathcal{V}_{n,p}$ (the elements of TL_n are scalars as far as the tensor product is concerned). Indeed, $\mathcal{V}_{n,p}\!\uparrow$ may alternatively be characterised as the quotient of the $\mathbb{C}$-tensor product $\mathsf{TL}_{n+1} \otimes_{\mathbb{C}} \mathcal{V}_{n,p}$ (with the TL_{n+1}-action $a\,(a' \otimes_{\mathbb{C}} z) = (aa') \otimes_{\mathbb{C}} z$ as before) by the submodule generated by the elements of the form $(ab \otimes_{\mathbb{C}} z - a \otimes_{\mathbb{C}} bz)$. This characterisation will shortly prove useful.

By Proposition 2.2, any monomial in the TL_{n+1}-generators u_i can be written in reverse Jones' normal form. In this form, the generator u_n appears at most once to the far left and may only be preceded by a string of the form $u_r u_{r+1} \cdots u_{n-1}$. We conclude that TL_{n+1} is spanned by monomials of the forms $u_r \cdots u_{n-1} u_n U$ and U', where U and U' are monomials in the TL_n-subalgebra. Thus, if D denotes a basis of $\mathcal{V}_{n,p}$, then $\mathcal{V}_{n,p}\!\uparrow$ is *spanned* by the set

$$\{\, \mathbf{1} \otimes d\ ,\ u_n \otimes d\ ,\ u_{n-1} u_n \otimes d\ , \ldots,\ u_1 \cdots u_{n-1} u_n \otimes d\ :\ d \in D \,\}. \tag{6.2}$$

Note that this spanning set is not usually a basis of $\mathcal{V}_{n,p}\!\uparrow$. For example, if d can be written as $u_{n-1} d'$ for some $d' \in \mathcal{V}_{n,p}$, then

$$u_{n-1} u_n \otimes d = u_{n-1} u_n \otimes u_{n-1} d' = u_{n-1} u_n u_{n-1} \otimes d' = u_{n-1} \otimes d' = \mathbf{1} \otimes u_{n-1} d' = \mathbf{1} \otimes d. \tag{6.3}$$

To obtain a basis, we generalise this computation. We will say that a link in a link state d is *simple* if it joins neighboring positions i and $i+1$. We then say that d has a simple link at i. Obviously, if $d = u_i d'$, where d and d' are link states, then d will have a simple link at i. The converse is the following:

Lemma 6.1. *If d is an (n,p)-link state with $n \geqslant 3$ which has a simple link at i, then d can always be expressed as $u_i d'$ with d' another (n,p)-link state. When $n = 2$, such an expression is valid if and only if $\beta \neq 0$.*

Proof. We construct d' explicitly to be the (n,p)-link state which is identical to d except that the simple link joining i and $i+1$ is swapped with whatever appears at $i-1$ (or $i+1$). Schematically,

$$d =$$



$$\Rightarrow \qquad d' =$$



$$. \tag{6.4}$$

It is clear that d' satisfies $u_i d' = d$ and likewise clear that this construction requires $n \geqslant 3$. The only remaining possibility then concerns the unique $(2,1)$-link state $d =$. This satisfies $d = \beta^{-1} u_1 d$ when $\beta \neq 0$, but when $\beta = 0$, u_1 acts as the zero operator on $\mathcal{V}_{2,1}$. ∎

This lemma allows us to reduce the number of elements in the set (6.2) while preserving its spanning property. But before describing this reduction, we introduce some more vocabulary. The (n,p)-link state d will be called *r-admissible*, $1 \leqslant r \leqslant n$, if d has no simple link at r or below, meaning at any i with $i \geqslant r$. It follows that all (n,p)-link states are n-admissible. Moreover, an element $u_r u_{r+1} \cdots u_n \otimes d \in \mathcal{V}_{n,p}\!\uparrow$ of (6.2) will be called *admissible* if d is r-admissible.

Lemma 6.2. *Let $n \geqslant 3$ and $0 \leqslant p \leqslant \lfloor n/2 \rfloor$. Let $u \in \mathsf{TL}_{n+1}$ be a word in the generators and d be an (n,p)-link state. Then, there exists an integer s, $1 \leqslant s \leqslant n+1$, and an element $e \in \mathcal{V}_{n,p}$ that is either 0 or an s-admissible (n,p)-link state, such that $u \otimes d = u_s u_{s+1} \cdots u_n \otimes e$ in $\mathcal{V}_{n,p}\!\uparrow$ (if $s = n+1$, then $u \otimes d = \mathbf{1} \otimes e$).*

Proof. We suppose that the word u is written in reverse Jones' normal form. If u does not contain u_n, then $u \otimes d = \mathbf{1} \otimes ud$ and the form suggested is obtained by choosing $s = n+1$ and $e = ud$. Suppose now that u is $u_r u_{r+1} \cdots u_n u'$ where u' is a word of TL_n and therefore $u \otimes d = u_r u_{r+1} \cdots u_n \otimes u'd$. If the element $u'd$ is zero in $\mathcal{V}_{n,p}$, then again $u_r u_{r+1} \cdots u_n \otimes e$ is of the desired form with $e = 0$. If $u'd$ is r-admissible, take $e = u'd$.

The only remaining case is when $u'd$ is not r-admissible. This means that there is a simple link at r or below. Let t be the position of the lowest simple link. Clearly $r \leqslant t \leqslant n-1$ and thus $u_t \in \mathsf{TL}_n$. Because $n \geqslant 3$, Lemma 6.1 ensures the existence of an (n,p)-link state f such that $u_t f = u'd$. Then,

$$\begin{aligned}
u_r u_{r+1} \cdots u_n \otimes u'd &= u_r \cdots u_n \otimes u_t f \\
&= u_r \cdots u_{t-1} u_t u_{t+1} u_t u_{t+2} \cdots u_n \otimes f \\
&= u_r \cdots u_{t-1} u_t u_{t+2} \cdots u_n \otimes f \\
&= u_{t+2} \cdots u_n \otimes u_r \cdots u_t f \\
&= u_{t+2} \cdots u_n \otimes u_r \cdots u_{t-1} u'd.
\end{aligned} \tag{6.5}$$

(The two extreme values of t lead to $u_{r+2}\cdots u_n\otimes u'd$ for $t=r$ and $\mathbf{1}\otimes u_r\cdots u_{n-2}u'd$ for $t=n-1$.) The configurations of $u'd$ and of $u_r\cdots u_{t-1}u'd$ are shown in the following diagrams:

If $u'd =$  , then $u_r\cdots u_{t-1}u'd =$  $=$  .

Note that in the lower box of $u'd$, marked by three dots, all positions are either defects or linked with positions in the two top boxes, since the lowest simple link is at t. (The two top boxes may have defects, links within and between them and, as just said, links with the bottom box.) The action of $u_r\cdots u_{t-1}$ is seen to exchange the simple link at t and the middle box. Since, after this exchange, the first position in the lower box is $t+2$, then the resulting link state is $(t+2)$-admissible. Clearly this is of the desired form if we set $s=t+2$ (and then $s\leqslant n+1$) and e is taken to be $u_r\cdots u_{t-1}u'd$. ■

Let $\mathcal{S}_{n,p}$ be the set consisting of all elements $\mathbf{1}\otimes d$ for any (n,p)-link state d and of elements $u_ru_{r+1}\cdots u_n\otimes d$ for all $1\leqslant r\leqslant n$ and all r-admissible (n,p)-link states d. Due to the previous lemma, $\mathcal{S}_{n,p}$ is a spanning set of $\mathcal{V}_{n,p}\!\uparrow$ for $n\geqslant 3$. (One can check that $\mathcal{S}_{2,p}$ spans $\mathcal{V}_{2,p}\!\uparrow$ except when $p=1$ and $\beta=0$.) As an example,



(6.6)

and $|\mathcal{S}_{3,1}|=5$. Our aim is to show that $\mathcal{S}_{n,p}$ is actually a basis of $\mathcal{V}_{n,p}\!\uparrow$. For this, we first introduce a linear map $d\mapsto \underline{d}$ between the vector spaces underlying $\mathcal{V}_{n,p}$ and $\mathcal{V}_{n+2,p+1}$. This is defined on the basis of (n,p)-link states by adding two points to each basis element and a simple link from $n+1$ to $n+2$:

$\underline{d} =$  . (6.7)

We now define

$$\Phi\colon \mathcal{V}_{n,p}\!\uparrow\ \to \mathcal{V}_{n+2,p+1}\!\downarrow, \qquad \Phi(u\otimes d)=u\underline{d} \qquad (u\in \mathsf{TL}_{n+1},\ d\in\mathcal{V}_{n,p}). \tag{6.8}$$

Our first concern is to check whether Φ is well-defined, that is, whether the images $\Phi(u\otimes d)$ and $\Phi(v\otimes e)$ are equal when $u\otimes d=v\otimes e$ for $u,v\in\mathsf{TL}_{n+1}$ and $d,e\in\mathcal{V}_{n,p}$. From the characterisation of induced modules as quotients of the $\mathbb{C}$-tensor product, it follows that Φ will be well-defined if $\Phi(uu'\otimes d)=\Phi(u\otimes u'd)$ for all $u\in\mathsf{TL}_{n+1}$, $u'\in\mathsf{TL}_n$ and $d\in\mathcal{V}_{n,p}$. But, this follows immediately from the associativity of the TL_{n+1}-action on $\mathcal{V}_{n+2,p+1}\!\downarrow$:

$$\Phi(uu'\otimes d)=(uu')\underline{d}=u(u'\underline{d})=u(\underline{u'd})=\Phi(u\otimes u'd). \tag{6.9}$$

We remark that the u' in $(u'\underline{d})$ should be interpreted as the image of $u' \in \mathsf{TL}_n$ in TL_{n+2} under the inclusion which adds two defects at the bottom.

The associativity of the TL_{n+1}-action also makes Φ a homomorphism of TL_{n+1}-modules (here is where the restriction is necessary):

$$u'\Phi(u \otimes d) = u'(u\underline{d}) = (u'u)\underline{d} = \Phi(u'u \otimes d) \qquad \text{for all } u,u' \in \mathsf{TL}_{n+1},\ d \in \mathcal{V}_{n,p}. \tag{6.10}$$

To prove that it is actually an isomorphism does not require much more effort.

Proposition 6.3.

(i) *For all* $n \geqslant 1$, $0 \leqslant p \leqslant \lfloor n/2 \rfloor$ *and* $\beta \in \mathbb{C}$, *the set* $\mathcal{S}_{n,p}$ *is a basis of* $\mathcal{V}_{n,p}\!\uparrow$, *except for* $\mathcal{V}_{2,1}\!\uparrow$ *at* $\beta = 0$ *which has instead* $\{\mathbf{1} \otimes$ $, u_2 \otimes$ $, u_1u_2 \otimes$ $\}$ *as a basis.*

(ii) *When* $(n,p) \neq (2,1)$ *or* $\beta \neq 0$, *the* TL_{n+1}*-modules* $\mathcal{V}_{n,p}\!\uparrow$ *and* $\mathcal{V}_{n+2,p+1}\!\downarrow$ *are isomorphic.*

Proof. The basis for $\mathcal{V}_{2,1}\!\uparrow$ is obtained by direct computation. For the general case, we show that the homomorphism Φ is both surjective and injective. The two statements *(i)* and *(ii)* are clear consequences.

Since the $(n+2,p+1)$-link states define a basis of $\mathcal{V}_{n+2,p+1}\!\downarrow$, the surjectivity of Φ will follow if every $(n+2,p+1)$-link state has a preimage in $\mathcal{S}_{n,p}$. Suppose that e is such a link state and note that e has at least one simple link. To construct $u \otimes d \in \mathcal{S}_{n,p}$ such that $\Phi(u \otimes d) = e$, let r be the position of the lowest simple link in e. Denote by d the (n,p)-link state obtained from e by deleting this lowest simple link. Since there are no simple links of e below r, the link d is r-admissible and therefore $u_ru_{r+1}\cdots u_n \otimes d = u \otimes d \in \mathcal{S}_{n,p}$. (Note that if $r = n+1$, then $u_ru_{r+1}\cdots u_n$ is understood to be $\mathbf{1}$.) It is now easy to verify that $\Phi(u_ru_{r+1}\cdots u_n \otimes d) = e$:

If $e =$ , then $d =$ 

and $\Phi(u_ru_{r+1}\cdots u_n \otimes d) = u_ru_{r+1}\cdots u_n\underline{d} =$  $=$  $= e.$ (6.11)

Therefore, every $(n+2,p+1)$-link state has a preimage in $\mathcal{S}_{n,p}$ and Φ is surjective.

The map Φ sends every element of $\mathcal{S}_{n,p}$ to an $(n+2,p+1)$-link state. Since the latter are linearly independent, the injectivity of Φ will be established if distinct elements of $\mathcal{S}_{n,p}$ have distinct images.

Suppose then that $w_1 = u_r u_{r+1} \cdots u_n \otimes d$ and $w_2 = u_s u_{s+1} \cdots u_n \otimes d'$ are elements of $\mathcal{S}_{n,p}$ with the same image. The link state d, being r-admissible, can be represented as above by two blocks, that marked by two dots being devoid of simple links. The image $\Phi(w_1)$ is then represented by the above diagram for e whose lowest simple link is at r. Similarly, the lowest simple link of $\Phi(w_2)$ must be at s. If $\Phi(w_1) = \Phi(w_2)$, then r and s must be equal. But then, the two link states $\Phi(w_1)$ and $\Phi(w_2)$ will coincide if and only if the upper boxes (marked by one dot) of d and d' coincide and similarly for their lower boxes (marked by two dots). This forces the original d and e to be equal and thus $w_1 = w_2$. ■

For completeness, we mention that as $\mathcal{V}_{2,1} \cong \mathcal{V}_{2,0}$ when $\beta = 0$, it follows that $\mathcal{V}_{2,1}\!\uparrow\, \cong \mathcal{V}_{3,1} \oplus \mathcal{V}_{3,0}$. However, Proposition 4.1 gives $\mathcal{V}_{4,2}\!\downarrow\, \cong \mathcal{V}_{3,1}$.

Because restricting a module does not change its dimension, the dimension of $\mathcal{V}_{n,p}\!\uparrow$ is that of $\mathcal{V}_{n+2,p+1}$. We therefore obtain:

Corollary 6.4. *The dimension of the induced module $\mathcal{V}_{n,p}\!\uparrow$ is*

$$\dim \mathcal{V}_{n,p}\!\uparrow\; = \begin{cases} 3 & \text{if } (n,p) = (2,1) \text{ and } \beta = 0, \\ d_{n+2,p+1} & \text{otherwise.} \end{cases} \tag{6.12}$$

Moreover, the structure of induced modules now follows, using the isomorphism Φ, from Proposition 4.1 and Corollary 4.2 which describe the structure of restricted ones.

Corollary 6.5. *When $(n,p) \neq (2,1)$ or $\beta \neq 0$, the sequence*

$$0 \longrightarrow \mathcal{V}_{n+1,p+1} \longrightarrow \mathcal{V}_{n,p}\!\uparrow\; \longrightarrow \mathcal{V}_{n+1,p} \longrightarrow 0 \tag{6.13}$$

is exact.

Corollary 6.6. *When (n,p) is non-critical, the exact sequence* (6.13) *splits, so we have (for $(n,p) \neq (2,1)$ or $\beta \neq 0$)*

$$\mathcal{V}_{n,p}\!\uparrow\; \cong \mathcal{V}_{n+1,p+1} \oplus \mathcal{V}_{n+1,p} \qquad \textit{(as } \mathsf{TL}_{n+1}\textit{-modules).} \tag{6.14}$$

Whether or not the exact sequence (6.13) splits, a submodule isomorphic to $\mathcal{V}_{n+1,p+1}$ is easily identified in $\mathcal{V}_{n,p}\!\uparrow$. In $\mathcal{V}_{n+2,p+1}\!\downarrow$, such a submodule is spanned by those $(n+2,p+1)$-link states which have a defect at position $n+2$ (see the proof of Proposition 4.1). Their images under Φ^{-1} have the form $u_r u_{r+1} \cdots u_n \otimes d$, with $r \leqslant n$ and d an (n,p)-link state with a defect at n. Then the injection $\mathcal{V}_{n+1,p+1} \to \mathcal{V}_{n,p}\!\uparrow$, call it α, of the exact sequence (6.13) sends $(n+1,p+1)$-link states to elements of $\mathcal{S}_{n,p}$ as follows: If d is an $(n+1,p+1)$-link state, then $\alpha(d) = u_r u_{r+1} \cdots u_n \otimes d'$, where the lowest simple link in d is at r and d' is the (n,p)-link state obtained from d by removing this simple link and adding a defect at position n.

The exact sequences in Equations (4.2) and (6.13) can be read off the Bratteli diagram quite easily. For restriction, the TL_{n-1}-modules $\mathcal{V}_{n-1,p}$ and $\mathcal{V}_{n-1,p-1}$ appearing in the exact sequence (4.2) for $\mathcal{V}_{n,p}\!\downarrow$ correspond to the entries immediately above, and to the left and right, respectively, of the entry corresponding to $\mathcal{V}_{n,p}$. For induction, we must instead look immediately below, and again to the left and right, to find the entries indicating the constituents of the exact sequence (6.13). We have learned that these exact sequences split when the module being restricted or induced is non-critical. The question of whether the sequences split when the module is critical will not be resolved until Section 8.

We conclude with an example showing that the induced module $\mathcal{V}_{n,p}\!\uparrow$ may be different to the direct sum $\mathcal{V}_{n+1,p+1} \oplus \mathcal{V}_{n+1,p}$ and even to the quotient $\mathcal{M}_{n+1,p}/\mathcal{M}_{n+1,p+2}$. These three TL_{n+1}-modules share the same exact sequence (6.13), the latter because Equations (3.2) and (3.3) give

$$\mathcal{V}_{n+1,p+1} = \frac{\mathcal{M}_{n+1,p+1}}{\mathcal{M}_{n+1,p+2}} \subseteq \frac{\mathcal{M}_{n+1,p}}{\mathcal{M}_{n+1,p+2}} \qquad \text{and} \qquad \frac{\mathcal{M}_{n+1,p}/\mathcal{M}_{n+1,p+2}}{\mathcal{M}_{n+1,p+1}/\mathcal{M}_{n+1,p+2}} \cong \frac{\mathcal{M}_{n+1,p}}{\mathcal{M}_{n+1,p+1}} = \mathcal{V}_{n+1,p}. \tag{6.15}$$

For our example, we take $q = e^{\mathrm{i}\pi/3}$, so that $\beta = 1$ and $(n,p) = (2,0)$ is critical. Then, one can check that the central element $F_3 \in \mathsf{TL}_3$, introduced in Appendix A, is represented on $\mathcal{V}_{2,0}\!\uparrow$ and $\mathcal{M}_3 = \mathcal{M}_{3,0}/\mathcal{M}_{3,2}$, with respective (ordered) bases $\Big\{u_2 \otimes$ $, u_1 u_2 \otimes$ $, \mathbf{1} \otimes$ $\Big\}$ and $\Big\{$, , $\Big\}$, by

$$\begin{pmatrix} -1 & 0 & 3 \\ 0 & -1 & -3 \\ 0 & 0 & -1 \end{pmatrix} \qquad \text{and} \qquad \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}. \tag{6.16}$$

Thus, this central element can be diagonalised on $\mathcal{M}_3$ (and on $\mathcal{V}_{3,1} \oplus \mathcal{V}_{3,0}$ by Proposition 3.8), but not on $\mathcal{V}_{2,0}\!\uparrow$.

7 The Irreducibility of the Radicals

The present section carries on the exploration, launched in Section 5, of the standard modules at roots of unity. The goal here is to prove that the radicals $\mathcal{R}_{n,p}$ of the standard modules $\mathcal{V}_{n,p}$ are either trivial (meaning $\mathcal{R}_{n,p} = \{0\}$) or irreducible. One can actually be more precise: Let q be a root of unity with ℓ the smallest positive integer such that $q^{2\ell} = 1$. We shall say that two pairs (n, p') and (n, p) with $0 < |p' - p| < \ell$ form a *symmetric pair* if (n, p') and (n, p) are non-critical and are located symmetrically on either side of the (single) critical line between them. The adjective *symmetric* will also be used when two objects, for example $\mathcal{V}_{n,p'}$ and $\mathcal{V}_{n,p}$, are labelled by a symmetric pair (n, p') and (n, p). The relevance of this concept is already apparent from previous sections. In particular, Proposition 5.3 equates the dimensions of the symmetric pair (with $p' > p$) $\mathcal{R}_{n,p'}$ and $\mathcal{L}_{n,p}$. In this section, we shall prove that the modules $\mathcal{R}_{n,p'}$ and $\mathcal{L}_{n,p}$ of such a symmetric pair are *isomorphic*. We remark that, as noted at the end of Section 4, every (n, p) is critical when $\beta = \pm 2$ or when $\beta = 0$ with n odd. It follows that there are no symmetric pairs in these cases. However, for all other β and n, such pairs exist whenever $n \geqslant \ell$.

A crucial role will be played in this section by the central element F_n of TL_n whose detailed properties may be found in Appendix A. The first part of this section shows that, even though F_n is central, its action is non-diagonalisable on certain indecomposable modules. The second part then uses this fact to construct an isomorphism between the symmetric pair $\mathcal{R}_{n,p'}$ and $\mathcal{L}_{n,p}$ (with $p' > p$). The inspiration here comes from the theory of staggered modules in logarithmic conformal field theory, see [20] for example. Though the irreducibility of the radical is well-known to experts, the only proof that we are aware of [17] relies upon some rather abstract category-theoretic analysis.[5] The last part describes completely the space of homomorphisms between standard modules.

Proposition A.2 states that F_n acts as a multiple of the identity on the standard modules $\mathcal{V}_{n,p}$. The next lemma provides a simple example where this action is non-diagonal. It will turn out to be key for what follows.

Lemma 7.1. *Let q be a root of unity other than* ± 1 *and let* (n, p) *be critical for this q (so* $n \neq 2p$*). If* z_p *denotes the* (n, p)*-link state which has p simple links at* $1, 3, \dots, 2p-1$*, then when* $F_{n+1}(\mathbf{1} \otimes z_p)$ *is expanded in the basis* $\mathcal{S}_{n,p}$ *of* $\mathcal{V}_{n,p}\!\uparrow$*, the coefficient of* $u_1 u_2 \cdots u_n \otimes z_p = u_{2p+1} u_{2p+2} \dots u_n \otimes z_p$ *does not vanish.*

Proof. We first study the case $p = 0$. Since $F_{n+1} \in \mathsf{TL}_{n+1}$, it can be written as a sum of words in reverse Jones' normal form, as in Equation (2.7). For such a word to act non-trivially on $\mathbf{1} \otimes z_0$, the rightmost u_j must be u_n. So only words with a single "flight" $u_i u_{i+1} \cdots u_n$ will contribute, and among these, only $u_1 u_2 \cdots u_n$ will lead to the term $u_1 u_2 \cdots u_n \otimes z_0$ that we are seeking. We therefore need to compute the coefficient of $u_1 u_2 \cdots u_n$ in F_{n+1}.

---

[5] Our proof is based on the existence of a non-zero homomorphism $\theta : \mathcal{V}_{n,p} \to \mathcal{V}_{n,p'}$ when the two modules form a symmetric pair with $p' > p$. Martin [2] has proved that $\ker(\theta)$ and $\operatorname{coker}(\theta)$ are irreducible. This follows from, but is weaker than, Theorem 7.2 below.

We first expand the two crossings of the top row of the diagram (A.2) defining F_{n+1}. The points marked by dots in the following diagrams need to be linked if they are to lead to the word $u_1u_2\cdots u_n$.

 $= -$  $+q^{-1}$  $-$  $+q$  . (7.1)

The last term will not contribute to $u_1u_2\cdots u_n$, so we obtain

 $\overset{\circ}{=} (-2+\beta q^{-1})$  , (7.2)

where the sign "$\overset{\circ}{=}$" indicates that the equality is restricted to the coefficient of $u_1u_2\cdots u_n$. We now expand the second row. Only one term of the crossing on the right may contribute, as the other term can be seen to close a link on the right vertical line. But then only one term of the crossing on the left contributes to $u_1u_2\cdots u_n$. A similar argument can be repeated for all rows but the last one and the expansion of all crossings but those of the bottom row gives

 $\overset{\circ}{=} (-2+\beta q^{-1})(-1)^{n-1}$  . (7.3)

Note that the top points on the left are now linked; those on the bottom right, marked by dots, still remain to be linked. Expanding the last row, we finally get that the coefficient of $u_1u_2\cdots u_n$ in F_{n+1} is

$$(-1)^{n-1}(-2+\beta q^{-1})(-2+\beta q) = (-1)^n(\beta^2-4). \tag{7.4}$$

Therefore, the coefficient of $u_1u_2\cdots u_n \otimes z_0$ in $F_{n+1}(\mathbf{1}\otimes z_0)$ vanishes only when $\beta = \pm 2$ ($q = \pm 1$).

We shall now assume that $p \geqslant 1$ and consider elements in $\mathcal{V}_{n,p}\!\uparrow$ of the form $v = \mathbf{1}\otimes v'$, where v' is constrained to be an (n,p)-link state with a simple link at 1. In other words,

$v = \mathbf{1}\otimes$  , (7.5)

where the box stands for any $(n, p-1)$-link state. We shall also make the hypothesis that $\beta \neq 0$ which allows us to use the identity

$$v = \beta^{-1}u_1 v \tag{7.6}$$

(the case $\beta = 0$ will be deferred until the end of the proof). We now expand the two crossings on the top right of F_{n+1}, extracting these crossings from the diagram representing F_{n+1} and using dots to mark the points at which the subdiagrams we have extracted are to be connected:





(where we have used the identity (7.6) on the first and last terms)



(7.7)

We now carry out the expansion of the two crossings on the left:



(7.8)

The first, third and last terms cancel and, by (7.6), the second becomes



(7.9)

Suppose now that the link state v' in $v = \mathbf{1} \otimes v'$ contains several consecutive simple links at the top. Then the above computation can be repeated for each of them with the result that


$F_{n+1}(\mathbf{1} \otimes z_p) =$
$2p$
F_{n-2p+1}
$(\mathbf{1} \otimes z_p),$
in $\mathcal{V}_{n,p}\!\uparrow.$

(7.10)

The expansion of the F_{n-2p+1} indicated at the bottom of this diagram will contain a term $u_{2p+1}u_{2p+2}\cdots u_n$ only if $n > 2p$. In that case, we may use the previous $p = 0$ computation to conclude that the coefficient of $u_1u_2\cdots u_n \otimes z_p$ in $F_{n+1}(\mathbf{1} \otimes z_p)$ is $(-1)^n(\beta^2 - 4)$ as before. Again this coefficient does not vanish under the hypotheses of the lemma.

The case $\beta = 0$, corresponding to $q = \pm i$, remains. In the link basis of the modules $\mathcal{V}_{n,p}$, the elements of the matrices representing the generators u_i are polynomials in β or, equivalently, in q

and q^{-1}. The matrices representing the u_i in the induced modules $\mathcal{V}_{n,p}\!\uparrow$ inherit this property if the basis $\mathcal{S}_{n,p}$ is used. Indeed, the action of u_i on an arbitrary member $u_r u_{r+1} \cdots u_n \otimes d$ of $\mathcal{S}_{n,p}$ is easily computed:

$$u_i\left(u_r \cdots u_n \otimes d\right) = \begin{cases} u_r \cdots u_n \otimes u_i d, & i < r-1 \\ u_{r-1} \cdots u_n \otimes d, & i = r-1 \\ \beta u_r \cdots u_n \otimes d, & i = r \\ u_i \cdots u_n \otimes u_r \cdots u_{i-2} d, & r+1 \leqslant i \leqslant n. \end{cases} \tag{7.11}$$

The right-hand sides are either elements of $\mathcal{S}_{n,p}$ or are equivalent to scalar multiples of such elements (using Lemma 6.2). The central element F_n, defined diagrammatically by (A.1) and (A.2), is a linear combination of words in the generators with weights that are also polynomials in q and q^{-1}. Therefore, the coefficients of $F_{n+1}(\mathbf{1} \otimes z_p)$ in the basis $\mathcal{S}_{n,p}$ of $\mathcal{V}_{n,p}\!\uparrow$ are polynomials in q and q^{-1}. Their actual values for a given β can be obtained by evaluation of these polynomials at the corresponding value of q. The coefficient of $u_1 u_2 \cdots u_n \otimes z_p$ in $F_{n+1}(\mathbf{1} \otimes z_p)$ when $\beta = 0$ is therefore $(-1)^n(\beta^2 - 4) = 4(-1)^{n+1}$ and is non-zero. ■

Equation (6.16) provides a simple example of this off-diagonal action — there, we considered F_3 acting on the indecomposable module $V_{2,0}\!\uparrow$ at $q = e^{\mathrm{i}\pi/3}$.

Proposition 3.6 has shown that the only homomorphism $\mathcal{V}_{n,p'} \to \mathcal{V}_{n,p}$ with $p' > p$ and $\langle \cdot, \cdot \rangle_{n,p}$ and $\langle \cdot, \cdot \rangle_{n,p'} \neq 0$ is the zero homomorphism. The above properties of the central element F_n allow us to construct a non-zero homomorphism $\mathcal{V}_{n,p} \to \mathcal{V}_{n,p'}$, $p' > p$, when the modules form a symmetric pair, and thereby reveal the structure of the radical $\mathcal{R}_{n,p}$.

Theorem 7.2. *The radical $\mathcal{R}_{n,p}$ is zero or irreducible and, if the pair $\mathcal{V}_{n,p}$ and $\mathcal{V}_{n,p'}$, with $p' > p$, is symmetric, then $\mathcal{R}_{n,p'} \cong \mathcal{L}_{n,p}$.*

Proof. If q is not a root of unity, these statements have already been proven. Indeed, the $\mathcal{V}_{n,p}$ are then irreducible (Proposition 4.5) and their radicals are trivial. The same is true when $q = \pm 1$ and when $q = \pm \mathrm{i}$ with n odd (Corollary 4.8). The theorem is therefore non-trivial only if q is a root of unity with either $\ell \geqslant 3$ or $\ell = 2$ and n even. We will therefore assume these conditions in what follows.

We first construct a non-zero homomorphism between the two standard modules $\mathcal{V}_{n+1,p}$ and $\mathcal{V}_{n+1,p+1}$, assuming that (n,p) is critical. The symmetric pair therefore consists of $(n+1,p)$ and $(n+1,p+1)$ and we note that $f_{n+1,p}$ and $f_{n+1,p+1}$ are equal. Let $\varphi \colon \mathcal{V}_{n,p}\!\uparrow \to \mathcal{V}_{n,p}\!\uparrow$ be the map obtained from left-multiplication by $F_{n+1} - f_{n+1,p}\mathbf{1} \in \mathsf{TL}_{n+1}$. Because this element is central, φ is a homomorphism. Moreover, it is non-zero by Lemma 7.1.

Consider now the exact sequence (6.13)

$$0 \longrightarrow \mathcal{V}_{n+1,p+1} \xrightarrow{\alpha} \mathcal{V}_{n,p}\!\uparrow \xrightarrow{\gamma} \mathcal{V}_{n+1,p} \longrightarrow 0 \tag{7.12}$$

(the homomorphism α was constructed explicitly after Corollary 6.6). Since φ acts as zero on $\mathcal{V}_{n+1,p+1}$ (Lemma A.2), we have $\operatorname{im}(\alpha) \subseteq \ker(\varphi)$. Similarly, $\gamma \circ \varphi = 0$ and therefore $\operatorname{im}(\varphi) \subseteq$

$\ker(\gamma) = \operatorname{im}(\alpha)$. Now, for any $w \in \mathcal{V}_{n+1,p}$, we can find $v \in \mathcal{V}_{n,p}\!\uparrow$ such that $\gamma(v) = w$. Note that any other $v' \in \mathcal{V}_{n,p}\!\uparrow$ with $\gamma(v') = w$ satisfies $v - v' \in \ker(\gamma) = \operatorname{im}(\alpha) \subseteq \ker(\varphi)$. It follows that the map $w \mapsto \varphi(v)$ is independent of the choice of preimage v. This map, in turn, is a TL_{n+1}-homomorphism from $\mathcal{V}_{n+1,p}$ into $\operatorname{im}(\alpha) \subset \mathcal{V}_{n,p}\!\uparrow$ since given any $u \in \mathsf{TL}_{n+1}$, we may choose uv as a preimage of uw and uw is mapped to $\varphi(uv) = u\varphi(v)$. Finally, let $i\colon \mathcal{V}_{n+1,p} \to \mathcal{V}_{n+1,p+1}$ be defined by $w \mapsto \left(\alpha^{-1} \circ \varphi\right)(v)$. This is the non-zero TL_{n+1}-homomorphism that we set out to construct. The conclusion is therefore that $\dim \operatorname{Hom}_{\mathsf{TL}_{n+1}}(\mathcal{V}_{n+1,p}, \mathcal{V}_{n+1,p+1}) \geqslant 1$, when (n,p) is critical.

Frobenius reciprocity (Proposition B.4) allows us to extend this result to arbitrary symmetric pairs. Assume again that (n,p) is critical. Then, the pairs $\mathcal{V}_{n+j,p}$ and $\mathcal{V}_{n+j,p+j}$ are symmetric for $1 \leqslant j < \ell$, where ℓ is the smallest positive integer such that $q^{2\ell} = 1$, and any symmetric pair is of this form for some j and critical (n,p). Suppose now that $j \in \{2, \cdots, \ell - 1\}$. We will justify each step of the following computation:

$$\begin{aligned}
\operatorname{Hom}_{\mathsf{TL}_{n+j}}(\mathcal{V}_{n+j,p}, \mathcal{V}_{n+j,p+j}) &= \operatorname{Hom}_{\mathsf{TL}_{n+j}}(\mathcal{V}_{n+j,p} \oplus \mathcal{V}_{n+j,p+1}, \mathcal{V}_{n+j,p+j}) \\
&= \operatorname{Hom}_{\mathsf{TL}_{n+j}}(\mathcal{V}_{n+j-1,p}\!\uparrow, \mathcal{V}_{n+j,p+j}) \\
&= \operatorname{Hom}_{\mathsf{TL}_{n+j-1}}(\mathcal{V}_{n+j-1,p}, \mathcal{V}_{n+j,p+j}\!\downarrow) \\
&= \operatorname{Hom}_{\mathsf{TL}_{n+j-1}}(\mathcal{V}_{n+j-1,p}, \mathcal{V}_{n+j-1,p+j} \oplus \mathcal{V}_{n+j-1,p+j-1}) \\
&= \operatorname{Hom}_{\mathsf{TL}_{n+j-1}}(\mathcal{V}_{n+j-1,p}, \mathcal{V}_{n+j-1,p+j-1}).
\end{aligned} \tag{7.13}$$

Recall that $f_{n,p}$ takes distinct values between any two adjacent critical lines and that $f_{n,p} \neq f_{n,p'}$ implies that $\operatorname{Hom}(\mathcal{V}_{n,p}, \mathcal{V}_{n,p'}) = \{0\}$. The first and last equalities in (7.13) follow from this observation and the usual properties of Hom. Since (n,p) is critical and $2 \leqslant j < \ell$, neither $(n+j-1,p)$ nor $(n+j,p+j)$ can be critical, so Corollary 6.6 explains the second equality and Corollary 4.2 the fourth. The third equality is Frobenius reciprocity. We therefore conclude that

$$\operatorname{Hom}_{\mathsf{TL}_{n+j}}(\mathcal{V}_{n+j,p}, \mathcal{V}_{n+j,p+j}) = \operatorname{Hom}_{\mathsf{TL}_{n+1}}(\mathcal{V}_{n+1,p}, \mathcal{V}_{n+1,p+1}) \neq \{0\}. \tag{7.14}$$

In other words, there exists a non-zero homomorphism $\mathcal{V}_{n,p} \to \mathcal{V}_{n,p'}$ between an arbitrary symmetric pair with $p' > p$.

We are now ready to prove the statement of the theorem. Let $f \in \operatorname{Hom}(\mathcal{V}_{n,p}, \mathcal{V}_{n,p'})$ be a non-zero homomorphism between a symmetric pair with $p' > p$. Then, $\ker(f)$ is a proper subset of $\mathcal{V}_{n,p}$ and must therefore be a subset of $\mathcal{R}_{n,p}$ by maximality of the radical. If f were surjective, then $\mathcal{V}_{n,p}/\ker(f) \cong \mathcal{V}_{n,p'}$ which would contradict Proposition 3.6. (Here, we must temporarily assume that $\beta \neq 0$ or $p' \neq n/2$ — see the end of the present paragraph.) Then, $\operatorname{im}(f)$ is a subset of $\mathcal{R}_{n,p'}$, again by maximality. If either of $\ker(f)$ or $\operatorname{im}(f)$ is a proper subset of the corresponding radical, then we would have

$$\dim \ker(f) < \dim \mathcal{R}_{n,p} \qquad \text{or} \qquad \dim \operatorname{im}(f) < \dim \mathcal{R}_{n,p'} \tag{7.15}$$

which gives

$$\dim \mathcal{V}_{n,p} = \dim \ker(f) + \dim \operatorname{im}(f) < \dim \mathcal{R}_{n,p} + \dim \mathcal{R}_{n,p'}. \tag{7.16}$$

But then, the dimension of the irreducible quotient $\mathcal{L}_{n,p} = \mathcal{V}_{n,p}/\mathcal{R}_{n,p}$ would satisfy

$$\dim \mathcal{L}_{n,p} = \dim \mathcal{V}_{n,p} - \dim \mathcal{R}_{n,p} < \dim \mathcal{R}_{n,p'}, \tag{7.17}$$

contradicting Proposition 5.3. This proves that $\ker(f) = \mathcal{R}_{n,p}$ and $\operatorname{im}(f) = \mathcal{R}_{n,p'}$. The first isomorphism theorem now says that $\mathcal{L}_{n,p} = \mathcal{V}_{n,p}/\ker(f)$ is isomorphic to $\operatorname{im}(f) = \mathcal{R}_{n,p'}$ if the pair (n,p) and (n,p'), $p' > p$, is symmetric. The case $\beta = 0$ and $2p' = n$ was set aside in this argument, but is in fact easier because in this case Corollary 4.9 has already established that $\mathcal{R}_{2p',p'} = \mathcal{V}_{2p',p'}$ is irreducible. Since f is non-zero, $\operatorname{im}(f) = \mathcal{R}_{2p',p'}$ (and $\dim\operatorname{im}(f) = \dim\mathcal{R}_{n,p'}$). The rest of the proof is identical.

This argument proves the irreducibility of every $\mathcal{R}_{n,p'}$ for which the image (n,p) of (n,p') by a reflection with respect to the critical line immediately to its right is well-defined (meaning $p \geqslant 0$). This test fails for some (n,p') on the right of the second-rightmost critical line. For example, Figure 2, drawn for $\ell = 4$, shows that $(4,0)$ and $(8,2)$ have no reflection with respect to the critical line on their right. For all pairs (n,p') without such reflection, Proposition 5.3 shows that the radicals attached to them are $\{0\}$. ■

Frobenius reciprocity was a key ingredient in the previous proof. It allowed us to deduce the existence of non-trivial homomorphisms between certain standard modules $\mathcal{V}_{n,p}$ from the explicit construction of such a homomorphism in the simplest case. This underlines also the usefulness of a central element, in the present case F_n, that acts non-diagonalisably. A natural question which remains is whether these non-trivial homomorphisms are the only ones. We conclude the section by settling this in the affirmative.

Theorem 7.3. *The dimension* $\dim\operatorname{Hom}(\mathcal{V}_{n,p},\mathcal{V}_{n,p'})$ *is* 1 *if* $p = p'$ *or if the two standard modules form a symmetric pair with* $p' > p$*. Moreover, there is a single exceptional case:* $\dim\operatorname{Hom}(\mathcal{V}_{2,1},\mathcal{V}_{2,0}) = 1$ *when* $\beta = 0$*. Otherwise,* $\dim\operatorname{Hom}(\mathcal{V}_{n,p},\mathcal{V}_{n,p'}) = 0$.

Proof. We first omit, for $\beta = 0$ and n even, the study of $\operatorname{Hom}(\mathcal{V}_{2p,p},\mathcal{V}_{2p,p'})$ and $\operatorname{Hom}(\mathcal{V}_{2p,p'},\mathcal{V}_{2p,p})$. This omission makes available all results from Section 3 that call for the bilinear form $\langle\cdot,\cdot\rangle$ to be non-zero. Then the homomorphisms $\mathcal{V}_{n,p} \to \mathcal{V}_{n,p'}$ have been determined by Proposition 3.6, in the case $p > p'$, and Proposition 3.8, in the case $p = p'$. We also know that $\operatorname{Hom}(\mathcal{V}_{n,p},\mathcal{V}_{n,p'}) = \{0\}$ when $f_{n,p} \neq f_{n,p'}$. The outstanding cases are therefore:

1) (n,p) and (n,p') are both critical with $p' > p$ and $f_{n,p} = f_{n,p'}$.
2) (n,p) and (n,p') form a symmetric pair with $p' > p$.
3) (n,p) and (n,p') are non-critical with $p' - p \geqslant \ell$ and $f_{n,p} = f_{n,p'}$.

Case (1) is easily dealt with: If two distinct (n,p) and (n,p') are critical, then $\mathcal{V}_{n,p}$ and $\mathcal{V}_{n,p'}$ are irreducible (Corollary 4.8) and it follows from Corollary 3.7 that $\operatorname{Hom}(\mathcal{V}_{n,p},\mathcal{V}_{n,p'}) = 0$. In case (2), we know from Theorem 7.2 that $\mathcal{R}_{n,p'} \cong \mathcal{L}_{n,p}$. Schur's lemma then implies that any two homomorphisms from $\mathcal{L}_{n,p}$ to $\mathcal{R}_{n,p'}$ are equal to one another up to a multiplicative constant. Choose two non-zero homomorphisms $f,g \in \operatorname{Hom}(\mathcal{V}_{n,p},\mathcal{V}_{n,p'})$. The argument used to prove Theorem 7.2 implies that we must necessarily have $\ker(f) = \ker(g) = \mathcal{R}_{n,p}$ and $\operatorname{im}(f) = \operatorname{im}(g) = \mathcal{R}_{n,p'}$. Both f and g then define homomorphisms $\hat{f},\hat{g}\colon \mathcal{L}_{n,p} \to \mathcal{R}_{n,p'}$ and so we learn that $\hat{f} = \mu\hat{g}$ for some $\mu \in \mathbb{C}$.

As

$$f(v) = \left(\hat{f} \circ \pi\right)(v) = \mu\left(\hat{g} \circ \pi\right)(v) = \mu g(v) \qquad \text{for all } v \in \mathcal{V}_{n,p}, \tag{7.18}$$

where π is the projection $\mathcal{V}_{n,p} \to \mathcal{L}_{n,p}$, we conclude that $f = \mu g$, hence that $\dim \operatorname{Hom}(\mathcal{V}_{n,p}, \mathcal{V}_{n,p'}) = 1$.

We therefore turn to case (3). Let (n, p'') be the reflection of (n, p') in the critical line immediately to its right, so that $\mathcal{V}_{n,p'}$ and $\mathcal{V}_{n,p''}$ form a symmetric pair with $p' > p'' > p$. Let $f \colon \mathcal{V}_{n,p} \to \mathcal{V}_{n,p'}$ be a non-zero homomorphism, so $\ker(f) \subseteq \mathcal{R}_{n,p}$. But, $\mathcal{R}_{n,p}$ is irreducible and therefore $\ker(f)$ is either $\{0\}$ or $\mathcal{R}_{n,p}$. Similarly, $\operatorname{im}(f)$ must be either $\mathcal{R}_{n,p'}$ or $\mathcal{V}_{n,p'}$. The first isomorphism theorem, together with Corollary 3.7, now leads to a contradiction in each of the four possible combinations. For example:

- If $\ker(f) = \{0\}$ and $\operatorname{im}(f) = \mathcal{V}_{n,p'}$, then $\mathcal{V}_{n,p} \cong \operatorname{im}(f) = \mathcal{V}_{n,p'}$ with $p \neq p'$, contradicting Corollary 3.7.
- If $\ker(f) = \{0\}$ and $\operatorname{im}(f) = \mathcal{R}_{n,p'}$, then $\mathcal{V}_{n,p} \cong \operatorname{im}(f) = \mathcal{R}_{n,p'} \cong \mathcal{L}_{n,p''}$. But, $\mathcal{V}_{n,p}$ being irreducible implies that $\mathcal{L}_{n,p} \cong \mathcal{L}_{n,p''}$ with $p \neq p''$, again contradicting Corollary 3.7.

The last two cases $\ker(f) = \mathcal{R}_{n,p}$, $\operatorname{im}(f) = \mathcal{V}_{n,p'}$ and $\ker(f) = \mathcal{R}_{n,p}$, $\operatorname{im}(f) = \mathcal{R}_{n,p'}$ are similar and are left to the reader. We conclude that $\operatorname{Hom}(\mathcal{V}_{n,p}, \mathcal{V}_{n,p'}) = \{0\}$ in all these cases.

It only remains to consider the omitted case involving $\mathcal{V}_{2p,p} = \mathcal{R}_{2p,p}$ at $\beta = 0$. In this case, every $(2p, p')$ is non-critical (see Figure 1) and shares the same F_{2p}-eigenvalue: $f_{2p,p'} = 0$. $\mathcal{V}_{2p,p'}$ and $\mathcal{V}_{2p,p''}$ therefore form a symmetric pair if and only if $|p' - p''| = 1$. Corollary 3.7 can be used to conclude that the $\mathcal{L}_{2p,p'}$ with $0 \leqslant p' \leqslant p-1$ are mutually non-isomorphic. Because $\mathcal{L}_{2p,p'} \cong \mathcal{R}_{2p,p'+1}$, the radicals $\mathcal{R}_{2p,p'}$ with $1 \leqslant p' \leqslant p$ are likewise mutually non-isomorphic. Finally, the $\mathcal{R}_{2p,p'} = \mathcal{L}_{2p,p'-1}$ are non-zero for $1 \leqslant p' < p$, so the only irreducible modules among the $\mathcal{V}_{2p,p'}$ are therefore $\mathcal{V}_{2p,p}$ (by Proposition 4.9) and $\mathcal{V}_{2p,0}$.

We now look for a non-zero homomorphism $f \colon \mathcal{V}_{2p,p} \to \mathcal{V}_{2p,p'}$ with $p' < p$. If such a homomorphism exists, then $\ker(f) = \{0\}$ and $\operatorname{im}(f) \cong \mathcal{V}_{2p,p}$ (because the latter is irreducible). It follows that either $\mathcal{V}_{2p,p} \cong \mathcal{R}_{2p,p'}$ or $\mathcal{V}_{2p,p} \cong \mathcal{V}_{2p,p'}$. The first possibility is excluded by the previous observation that the radicals $\mathcal{R}_{2p,p'}$, $1 \leqslant p' \leqslant p$, are mutually distinct as well as recalling that $\mathcal{R}_{2p,p} = \mathcal{V}_{2p,p} \neq \{0\} = \mathcal{R}_{2p,0}$. We are therefore left with the second possibility, that $\mathcal{V}_{2p,p'}$ is itself irreducible, which can only happen when $p' = 0$. But, $\dim \mathcal{V}_{2p,0} = 1$ only coincides with $\dim \mathcal{V}_{2p,p}$ when $p = 1$. So we arrive at the exceptional case $\operatorname{Hom}(\mathcal{V}_{2,1}, \mathcal{V}_{2,0}) \cong \mathbb{C}$ (the existence of such a non-zero homomorphism was already established after Corollary 3.7).

Finally, the pair $\mathcal{V}_{2p,p-1}$ and $\mathcal{V}_{2p,p}$ is symmetric and the same arguments that proved (2) above lead to $\dim \operatorname{Hom}(\mathcal{V}_{2p,p-1}, \mathcal{V}_{2p,p}) = 1$. The proof that $\dim \operatorname{Hom}(\mathcal{V}_{2p,p'}, \mathcal{V}_{2p,p}) = 0$ for $p' \leqslant p-2$ mimics that of (3) with the simplification that now $\operatorname{im}(f) = \mathcal{V}_{2p,p}$ since $\mathcal{V}_{2p,p}$ is irreducible, hence there are only two cases to consider instead of four. ∎

Since the only proper submodule of $\mathcal{V}_{n,p}$ is the radical $\mathcal{R}_{n,p}$ (and when $\beta = 0$, the exceptional case $\mathcal{V}_{2p,p}$ is irreducible), the only quotients obtained from $\mathcal{V}_{n,p}$ are the modules $\{0\}$, $\mathcal{V}_{n,p}/\mathcal{R}_{n,p} \cong \mathcal{L}_{n,p}$ and $\mathcal{V}_{n,p}$ itself. Of course some of these may coincide. Therefore:

Corollary 7.4. *Every quotient of a standard module is indecomposable.*

8 Principal Indecomposable Modules

In this section, we describe a concrete construction, for q a root of unity, of the *principal indecomposable* modules as submodules of certain induced modules. These are the indecomposable direct summands of the Temperley-Lieb algebra when one treats it as a module by letting it act on itself by left-multiplication (this is just the regular representation). It follows immediately that principal indecomposables are canonical examples of *projective* modules. Indeed, they are precisely the indecomposable projective modules. Moreover, a standard fact about them [21] is that there is a bijective correspondence between principal indecomposables and irreducibles given by quotienting the former by its (unique) maximal proper submodule (its radical). We will therefore denote a principal indecomposable module by $\mathcal{P}_{n,p}$, understanding that its irreducible quotient is $\mathcal{L}_{n,p}$.[6] The properties of projective modules needed for this section are reviewed in Appendix B.

Note that when the algebra is semisimple, indecomposability implies irreducibility, so

$$\mathcal{P}_{n,p} = \mathcal{V}_{n,p} = \mathcal{L}_{n,p} \qquad (\mathsf{TL}_n \text{ semisimple}). \tag{8.1}$$

We remark that because TL_n is semisimple for generic β (and its dimension is obviously independent of β), it follows from Theorems B.1 and B.3 that

$$\sum_i \dim \mathcal{L}_i \dim \mathcal{P}_i = \dim \mathsf{TL}_n = \sum_{p=0}^{\lfloor n/2 \rfloor} \dim \mathcal{V}_{n,p} \dim \mathcal{V}_{n,p}, \tag{8.2}$$

where the sum on the left-hand side is over a *complete set* of pairwise non-isomorphic irreducibles $\mathcal{L}_i$ and the corresponding principal indecomposables $\mathcal{P}_i$. We recall from Corollary 4.6 that the sum on the right-hand side corresponds to a complete set of non-isomorphic irreducibles, when q is not a root of unity. At this point, we know that the $\mathcal{L}_{n,p}$ are pairwise non-isomorphic, but we are not assured that they form a complete set when q is a root of unity, excluding $\mathcal{L}_{2p,p} = \{0\}$ from the set when $\beta = 0$. This completeness will be deduced as a corollary of the principal indecomposable analysis.

We will first show how one can concretely construct the principal indecomposables with the aid of a detailed example. This serves to nicely illustrate the salient features of the general discussion (which is obscured somewhat by the necessary induction arguments). So let us take $q = e^{\mathrm{i}\pi/4}$ ($\beta = \sqrt{2}$), noting that criticality corresponds to $n - 2p + 1$ being a multiple of $\ell = 4$. It may be helpful to recall that the dimensions of the irreducible modules for this β and small n were organized in a Bratteli diagram in Figure 2.

Consider first the rather trivial algebra TL_1. Since it is one-dimensional and spanned by the unit $\mathbf{1}$, its left regular representation is isomorphic to its only standard module $\mathcal{V}_{1,0}$. Thus,

$$\mathsf{TL}_1 = \mathcal{P}_{1,0} \cong \mathcal{V}_{1,0}. \tag{8.3}$$

---

[6] In the exceptional case, $\beta = 0$ and $n = 2p$, we recall that $\mathcal{L}_{2p,p} = \{0\}$ and the irreducible quotient of $\mathcal{V}_{2p,p}$ is in fact $\mathcal{V}_{2p,p} = \mathcal{R}_{2p,p} \cong \mathcal{L}_{2p,p-1}$. There is therefore no need to define a non-trivial principal indecomposable $\mathcal{P}_{2p,p}$ when $\beta = 0$.

Now apply the induced module construction to get $\mathsf{TL}_2 \otimes_{\mathsf{TL}_1} \mathsf{TL}_1 \cong \mathcal{V}_{1,0}\!\uparrow$ as TL_2-modules. Since $\mathsf{TL}_{n+1} \otimes_{\mathsf{TL}_n} \mathsf{TL}_n$ is obviously isomorphic to TL_{n+1} (just send $a \otimes b = ab \otimes \mathbf{1}$ to ab), we obtain

$$\mathsf{TL}_2 \cong \mathcal{V}_{1,0}\!\uparrow\ \cong \mathcal{V}_{2,1} \oplus \mathcal{V}_{2,0}. \tag{8.4}$$

Here, we have applied Corollary 6.6, using the fact that $(1,0)$ is not critical, because $\beta \neq 0$. Note that both $\mathcal{V}_{2,1}$ and $\mathcal{V}_{2,0}$ are indecomposable, by Proposition 3.3, and are direct summands of the left regular representation. They are therefore principal indecomposable modules: $\mathcal{P}_{2,1} \cong \mathcal{V}_{2,1}$ and $\mathcal{P}_{2,0} \cong \mathcal{V}_{2,0}$.

For $n = 3$, we similarly apply the induced module construction to Equation (8.4), obtaining

$$\mathsf{TL}_3 \cong (\mathsf{TL}_3 \otimes_{\mathsf{TL}_2} \mathcal{V}_{2,1}) \oplus (\mathsf{TL}_3 \otimes_{\mathsf{TL}_2} \mathcal{V}_{2,0}) \cong \mathcal{V}_{2,1}\!\uparrow\ \oplus \mathcal{V}_{2,0}\!\uparrow\ , \tag{8.5}$$

since tensor product distributes over direct sums. Now, $(2,0)$ is only critical for $\beta = \pm 1$, hence Corollary 6.6 applies, giving $\mathcal{V}_{2,0}\!\uparrow\ \cong \mathcal{V}_{3,1} \oplus \mathcal{V}_{3,0}$. Since $\mathcal{V}_{2,1}\!\uparrow\ \cong \mathcal{V}_{3,1}$, we conclude that $\mathcal{P}_{3,1} \cong \mathcal{V}_{3,1}$ and $\mathcal{P}_{3,0} \cong \mathcal{V}_{3,0}$. Note that this implies that $\mathsf{TL}_3 = 2\,\mathcal{P}_{3,1} \oplus \mathcal{P}_{3,0}$, consistent with Theorem B.3 as $\dim \mathcal{L}_{3,1} = 2$ and $\dim \mathcal{L}_{3,0} = 1$.

We pause to remark that for all n considered thus far, we have demonstrated the decomposition

$$\mathsf{TL}_n \cong \bigoplus_{p=0}^{\lfloor n/2 \rfloor} (\dim \mathcal{L}_{n,p})\, \mathcal{P}_{n,p}. \tag{8.6}$$

We may therefore deduce from Theorem B.3 that the set of $\mathcal{L}_{n,p}$ with $0 \leqslant p \leqslant \lfloor n/2 \rfloor$ constitutes a complete set of pairwise non-isomorphic irreducibles, at least for $n \leqslant 3$ and $\beta = \sqrt{2}$.

At some point, criticality must enter the fray. For $\beta = \sqrt{2}$, this first occurs for $n = 4$. Inducing our decomposition of TL_3 and analysing as above, we easily conclude that

$$\mathsf{TL}_4 \cong 2\,\mathcal{V}_{4,2} \oplus 2\,\mathcal{V}_{4,1} \oplus \mathcal{V}_{3,0}\!\uparrow\ , \tag{8.7}$$

hence that $\mathcal{P}_{4,2} \cong \mathcal{V}_{4,2}$ and $\mathcal{P}_{4,1} \cong \mathcal{V}_{4,1}$. Theorem B.3 requires that

$$\mathsf{TL}_4 \cong 2\,\mathcal{P}_{4,2} \oplus 2\,\mathcal{P}_{4,1} \oplus \mathcal{P}_{4,0} \oplus \cdots , \tag{8.8}$$

where the "$\oplus \cdots$" admits that there may be additional unknown principal indecomposables beyond the $\mathcal{P}_{4,p}$.[7] Comparing gives $\mathcal{V}_{3,0}\!\uparrow\ \cong \mathcal{P}_{4,0} \oplus \cdots$. Now, Corollary 6.6 does not help us to simplify $\mathcal{V}_{3,0}\!\uparrow$ because $(3,0)$ is critical. Instead, Corollary 6.5 gives the exact sequence

$$0 \longrightarrow \mathcal{V}_{4,1} \longrightarrow \mathcal{V}_{3,0}\!\uparrow\ \longrightarrow \mathcal{V}_{4,0} \longrightarrow 0. \tag{8.9}$$

We ask if this sequence splits. The next lemma shows that, if it does, then $\mathcal{V}_{3,0}\!\uparrow$ must split as $\mathcal{V}_{4,1} \oplus \mathcal{V}_{4,0}$. But then, TL_4 would have $\mathcal{P}_{4,1} \cong \mathcal{V}_{4,1}$ appearing as a summand with multiplicity 3, which is greater than $\dim \mathcal{L}_{4,1} = 2$, contradicting Theorem B.3. This shows that (8.9) does not split, so $\mathcal{V}_{3,0}\!\uparrow$ is indecomposable.

Lemma 8.1. *If* $0 \to A \to B \to C \to 0$ *is a short exact sequence of modules in which* A *and* C *are indecomposable, but* B *is decomposable, then the sequence splits and* $B \cong A \oplus C$.

---

[7] Such additional principal indecomposables would correspond to additional irreducibles. The existence of these would therefore amount to a negative answer to the question of the completeness of the set of $\mathcal{L}_{4,p}$.

Proof. Let $\imath$ be the inclusion of A into B and suppose that $B = B_1 \oplus B_2$ is a non-trivial decomposition. Then, $A \cong (\imath(A) \cap B_1) \oplus (\imath(A) \cap B_2)$, so either $\imath(A) \subseteq B_1$ or $\imath(A) \subseteq B_2$ by indecomposability. Say $\imath(A) \subseteq B_1$ without loss of generality. Then,

$$C \cong \frac{B_1 \oplus B_2}{\imath(A)} = \frac{B_1}{\imath(A)} \oplus B_2, \tag{8.10}$$

so $C \cong B_1/\imath(A)$ or $C \cong B_2$, again by indecomposability. But the first option makes $0 \to A \to B_1 \to C \to 0$ exact, whence $B_2 = \{0\}$ (contradiction). Thus, $C \cong B_2$ and hence $A \cong B_1$. ■

Since $\mathcal{L}_{4,0}$ is a quotient of $\mathcal{V}_{3,0}\!\uparrow$, we may therefore identify $\mathcal{P}_{4,0}$ as $\mathcal{V}_{3,0}\!\uparrow$. This means that there are no additional principal indecomposables to uncover, hence that the $\mathcal{L}_{4,p}$ furnish a complete set of pairwise non-isomorphic irreducibles. Indeed, one can check that $\mathsf{TL}_4 \cong 2\,\mathcal{P}_{4,2} \oplus 2\,\mathcal{P}_{4,1} \oplus \mathcal{P}_{4,0}$ is in perfect accord with Theorem B.3 since the dimensions of the irreducible $\mathcal{L}_{4,2}$, $\mathcal{L}_{4,1}$ and $\mathcal{L}_{4,0}$ are 2, 2 and 1, respectively (see Figure 2).

Continuing, we induce our $n = 4$ decomposition and argue as above to get

$$\mathsf{TL}_5 \cong 4\,\mathcal{P}_{5,2} \oplus 2\,\mathcal{P}_{5,1} \oplus \mathcal{P}_{4,0}\!\uparrow, \tag{8.11}$$

with $\mathcal{P}_{5,2} \cong \mathcal{V}_{5,2}$ and $\mathcal{P}_{5,1} \cong \mathcal{V}_{5,1}$. As $\dim \mathcal{L}_{5,2} = \dim \mathcal{L}_{5,1} = 4$ and $\dim \mathcal{L}_{5,0} = 1$, Theorem B.3 yields the decomposition

$$\mathcal{P}_{4,0}\!\uparrow \cong 2\,\mathcal{P}_{5,1} \oplus \mathcal{P}_{5,0} \oplus \cdots, \tag{8.12}$$

where we again allow the possibility of additional unknown principal indecomposables. We therefore need to analyse the structure of $\mathcal{P}_{4,0}\!\uparrow$. For this, we induce each module of the exact sequence (8.9). Because induction is right-exact (Proposition B.5), we obtain the exact sequence

$$\mathcal{V}_{4,1}\!\uparrow \cong \mathcal{V}_{5,2} \oplus \mathcal{V}_{5,1} \longrightarrow \mathcal{P}_{4,0}\!\uparrow \longrightarrow \mathcal{V}_{4,0}\!\uparrow \cong \mathcal{V}_{5,1} \oplus \mathcal{V}_{5,0} \longrightarrow 0 \tag{8.13}$$

which we have simplified using Corollary 6.6. Note that the central element F_5 has eigenvalue 0 on $\mathcal{V}_{5,2}$ and $\mathcal{V}_{5,0}$, but eigenvalue -2 on $\mathcal{V}_{5,1}$. Any homomorphism $A \to B$ will map generalised eigenspaces of A into generalised eigenspaces of B with the same eigenvalue f of F_n. It follows that $\mathcal{P}_{4,0}\!\uparrow$ decomposes into projectives as $\mathcal{P} \oplus \mathcal{P}'$ (see Proposition B.6) along the generalised eigenspaces of F_5 and that the exact sequence (8.13) is equivalent to the exactness of

$$\mathcal{V}_{5,1} \xrightarrow{\imath} \mathcal{P} \longrightarrow \mathcal{V}_{5,1} \longrightarrow 0 \qquad \text{and} \qquad \mathcal{V}_{5,2} \xrightarrow{\imath'} \mathcal{P}' \longrightarrow \mathcal{V}_{5,0} \longrightarrow 0. \tag{8.14}$$

From (8.12), we deduce that $\mathcal{P} = \mathcal{P}_{5,1} \oplus \mathcal{P}_{5,1}$ (hence $\ker \imath = \{0\}$) and that $\mathcal{P}' = \mathcal{P}_{5,0} \oplus \cdots$. It only remains to identify the submodule $\mathcal{W} = \mathcal{V}_{5,2}/\ker \imath'$ which makes the following sequence exact:

$$0 \longrightarrow \mathcal{W} \longrightarrow \mathcal{P}' \longrightarrow \mathcal{V}_{5,0} \longrightarrow 0. \tag{8.15}$$

Suppose first that (8.15) splits. Then, the projectivity of $\mathcal{P}'$ implies that $\mathcal{W}$ will be projective. But, $\mathcal{W}$ is then a projective quotient of the indecomposable $\mathcal{V}_{5,2}$, hence we must have $\mathcal{W} \cong \mathcal{V}_{5,2}$ or $\mathcal{W} = \{0\}$. The former possibility contradicts the multiplicity of $\mathcal{P}_{5,2} \cong \mathcal{V}_{5,2}$ in TL_5 being 4. So, $\mathcal{W} = \{0\}$ and $\mathcal{P}' \cong \mathcal{V}_{5,0}$ is indecomposable. If we instead suppose that (8.15) does not split, then Lemma 8.1 also gives us the indecomposability of $\mathcal{P}'$ because the only possibilities for $\mathcal{W}$ are indecomposable by Corollary 7.4. Either way, we may identify $\mathcal{P}'$ with $\mathcal{P}_{5,0}$ and conclude that the

$\mathcal{L}_{5,p}$ constitute a complete set of pairwise non-isomorphic irreducibles. To finally determine $\mathcal{W}$, we may now appeal to Equation (8.2), which gives

$$4\dim\mathcal{P}_{5,2}+4\dim\mathcal{P}_{5,1}+\dim\mathcal{P}_{5,0}=5\dim\mathcal{V}_{5,2}+4\dim\mathcal{V}_{5,1}+\dim\mathcal{V}_{5,0} \tag{8.16a}$$

$$\Rightarrow \qquad \dim\mathcal{P}_{5,0}=\dim\mathcal{V}_{5,2}+\dim\mathcal{V}_{5,0} \qquad \Rightarrow \qquad \mathcal{W}=\mathcal{V}_{5,2}. \tag{8.16b}$$

Hence, we have obtained the non-split exact sequence $0\to\mathcal{V}_{5,2}\to\mathcal{P}_{5,0}\to\mathcal{V}_{5,0}\to 0$. This completes the analysis of the principal indecomposables for $n=5$.

One can continue to analyse this example for higher n with few further difficulties. However, we have seen enough tricks by now to understand the principal indecomposables in general.

Theorem 8.2. *Let q be a root of unity and let ℓ be the minimal positive integer satisfying $q^{2\ell}=1$. Define $k(n,p)\in\mathbb{N}$ and $r(n,p)\in\{1,\ldots,\ell-1,\ell\}$ by $n-2p+1=k(n,p)\ell+r(n,p)$. Then, the principal indecomposables $\mathcal{P}_{n,p}$ of TL_n are identified as follows:*

- *If $r(n,p)=\ell$ (so (n,p) is critical), $\mathcal{P}_{n,p}\cong\mathcal{V}_{n,p}$.*
- *If $k(n,p)=0$ (so $n-2p+1\leqslant\ell$) and $\beta\neq 0$, then $\mathcal{P}_{n,p}\cong\mathcal{V}_{n,p}$.*
- *If $k(n,p)>0$ and $r(n,p)\neq\ell$, then $\mathcal{P}_{n,p}$ is the direct summand of the $r(n,p)$-fold induced module*

$$\mathcal{V}_{n-r(n,p),p}\ \underbrace{\uparrow\cdots\uparrow}_{r(n,p)\ \text{times}}$$

  consisting of the generalised eigenspace, under the action of F_n, whose generalised eigenvalue is $f_{n,p}=q^{n-2p+1}+q^{-(n-2p+1)}$. Furthermore, there is a non-split exact sequence

$$0\longrightarrow\mathcal{V}_{n,p+r(n,p)}\longrightarrow\mathcal{P}_{n,p}\longrightarrow\mathcal{V}_{n,p}\longrightarrow 0. \tag{8.17}$$

Moreover, $\{\mathcal{L}_{n,p}\ :\ p=0,1,\ldots,\lfloor n/2\rfloor\}$ is a complete set of pairwise non-isomorphic irreducibles, except when n is even and $\beta=0$. In this latter case, the range must be restricted to $p=0,1,\ldots,n/2-1$.

The proof is by induction on n and $k(n,p)$. As in the $\ell=4$ example detailed above, the key is to note that applying the induced module construction to a projective TL_{n-1}-module results in a projective TL_n-module. The labour mostly concerns keeping track of the multiplicities with which these projectives appear. Before commencing the induction arguments, it is convenient to deal with the exceptional cases $\beta=0,\pm2$. Indeed, the above statement makes it clear that the case $\beta=0$ and n even is special, since there is now no principal indecomposable $\mathcal{P}_{2p,p}$, the number of distinct principal indecomposables being then one less.

Proposition 8.3. *When either $\beta=\pm2$ ($\ell=1$) or n is odd and $\beta=0$ ($\ell=2$), the principal indecomposables of TL_n may be identified as $\mathcal{P}_{n,p}\cong\mathcal{V}_{n,p}\cong\mathcal{L}_{n,p}$, for $p=0,1,\ldots,n/2$. Moreover, the $\mathcal{L}_{n,p}$ form a complete set of pairwise non-isomorphic irreducibles for $0\leqslant p\leqslant n/2$.*

When n is even and $\beta=0$, we have instead $\mathcal{P}_{n,p}\cong\mathcal{V}_{n-1,p}\!\uparrow$, for $p=0,1,\ldots,n/2-1$. The $\mathcal{L}_{n,p}$ therefore form a complete set of pairwise non-isomorphic irreducibles for $0\leqslant p\leqslant n/2-1$.

Proof. When $\beta = \pm 2$ or n is odd and $\beta = 0$, TL_n is semisimple (as remarked after Corollary 4.8), hence $\mathcal{P}_{n,p} \cong \mathcal{V}_{n,p} = \mathcal{L}_{n,p}$ for all p. So, assume that n is even and $\beta = 0$. Then, $\ell = 2$ and the semisimplicity of TL_{n-1} gives

$$\mathsf{TL}_{n-1} \cong \bigoplus_{p=0}^{n/2-1} (\dim \mathcal{L}_{n-1,p})\, \mathcal{V}_{n-1,p}, \tag{8.18}$$

by Theorem B.1, hence

$$\mathsf{TL}_n \cong \bigoplus_{p=0}^{n/2-1} (\dim \mathcal{L}_{n-1,p})\, \mathcal{V}_{n-1,p}\!\uparrow\; = \bigoplus_{p=0}^{n/2-1} (\dim \mathcal{L}_{n,p})\, \mathcal{V}_{n-1,p}\!\uparrow. \tag{8.19}$$

Here, we have used Corollary 5.2, noting that $r(n,p) = 1 = \ell - 1$.

If we can show that the $\mathcal{V}_{n-1,p}\!\uparrow$ are indecomposable, then we may identify them with the $\mathcal{P}_{n,p}$. Theorem B.3 would then give the completeness of the corresponding $\mathcal{L}_{n,p}$. Now, this indecomposability is equivalent to

$$0 \longrightarrow \mathcal{V}_{n,p+1} \longrightarrow \mathcal{V}_{n-1,p}\!\uparrow\; \longrightarrow \mathcal{V}_{n,p} \longrightarrow 0 \tag{8.20}$$

not splitting, by Lemma 8.1. However, if this exact sequence splits, then $\mathcal{V}_{n,p+1}$ would be a direct summand of TL_n, hence it would be projective. The corresponding exact sequence for $\mathcal{V}_{n-1,p+1}\!\uparrow$ would then be guaranteed to split by statement *(ii)* of Proposition B.7. We therefore conclude that all the $\mathcal{V}_{n-1,p}\!\uparrow$ will be indecomposable, provided that the exact sequence for $\mathcal{V}_{n-1,n/2-1}\!\uparrow$ does not split.

Suppose then that $\mathcal{V}_{n-1,n/2-1}\!\uparrow\; \cong \mathcal{V}_{n,n/2} \oplus \mathcal{V}_{n,n/2-1}$. It follows that $\mathcal{V}_{n,n/2-1}$ and $\mathcal{V}_{n,n/2}$ would both be principal indecomposables of TL_n. Since $\mathcal{V}_{n,n/2} = \mathcal{R}_{n,n/2} \cong \mathcal{L}_{n,n/2-1}$ by Theorem 7.2, then both would be indecomposables with the same irreducible quotient $\mathcal{L}_{n,n/2-1}$ and should therefore coincide by Theorem B.3: $\mathcal{P}_{n,n/2-1} \cong \mathcal{V}_{n,n/2-1} \cong \mathcal{V}_{n,n/2}$. But, $\mathcal{V}_{n-1,n/2-1}\!\uparrow$ appears in TL_n with multiplicity $\dim \mathcal{L}_{n,n/2-1}$, by (8.19), hence this splitting would mean that $\mathcal{P}_{n,n/2-1}$ appears with multiplicity (at least) $2 \dim \mathcal{L}_{n,n/2-1}$. Since this contradicts Theorem B.3, the exact sequence for $\mathcal{V}_{n-1,n/2-1}\!\uparrow$ cannot split and the proof is complete. ■

As a preamble to the proof of Theorem 8.2, we gather some general facts concerning the result of inducing the principal indecomposables. Fix n and assume that the statement of the theorem is true for the principal indecomposables $\mathcal{P}_{n-1,p}$ and their irreducible quotients $\mathcal{L}_{n-1,p}$. We must consider two (non-exclusive) cases:

- *(i)* $\mathcal{P}_{n-1,p} \cong \mathcal{V}_{n-1,p}$, and
- *(ii)* $(n-1,p)$ is non-critical and there is a non-split exact sequence $0 \to \mathcal{V}_{n-1,p+r(n-1,p)} \to \mathcal{P}_{n-1,p} \to \mathcal{V}_{n-1,p} \to 0$.

In case *(i)*, $\mathcal{P}_{n-1,p}\!\uparrow$ is a projective TL_n-module with exact sequence

$$0 \longrightarrow \mathcal{V}_{n,p+1} \longrightarrow \mathcal{P}_{n-1,p}\!\uparrow\; \longrightarrow \mathcal{V}_{n,p} \longrightarrow 0. \tag{8.21}$$

If this splits, we identify $\mathcal{P}_{n,p+1} \cong \mathcal{V}_{n,p+1}$ and $\mathcal{P}_{n,p} \cong \mathcal{V}_{n,p}$, using the indecomposability of the standard modules (Proposition 3.3). If this does not split, then we instead find that $\mathcal{P}_{n,p} \cong \mathcal{P}_{n-1,p}\!\uparrow$. In case *(ii)*, we use the right-exactness of the induced module construction, the non-criticality of $(n-1,p)$ and the generalised eigenspaces of F_n to conclude that we have exact sequences

$$\mathcal{V}_{n,p+r(n,p)} \longrightarrow \mathcal{P} \longrightarrow \mathcal{V}_{n,p} \longrightarrow 0 \qquad \text{and} \qquad \mathcal{V}_{n,p+r(n,p)-1} \longrightarrow \mathcal{P}' \longrightarrow \mathcal{V}_{n,p+1} \longrightarrow 0, \qquad (8.22)$$

where $\mathcal{P}$ and $\mathcal{P}'$ are projectives satisfying $\mathcal{P} \oplus \mathcal{P}' = \mathcal{P}_{n-1,p}\!\uparrow$. If the first splits, then $\mathcal{P}_{n,p} \cong \mathcal{V}_{n,p}$ and $\mathcal{P}_{n,p+r(n,p)}$ will be a quotient of $\mathcal{V}_{n,p+r(n,p)}$ which, by Corollary 7.4, is indecomposable. If it does not, then we identify only $\mathcal{P}_{n,p}$. The same argument for the second exact sequence gives $\mathcal{P}_{n,p-1}$ and, perhaps, $\mathcal{P}_{n,p+r(n,p)-1}$.

This not only shows how one can identify principal indecomposables for TL_n from those of TL_{n-1}, it also indicates *which* TL_{n-1} principal indecomposables might produce a given TL_n principal indecomposable upon inducing (and decomposing). In particular, $\mathcal{P}_{n,p}$ can only be induced from

$$\mathcal{P}_{n-1,p}, \quad \mathcal{P}_{n-1,p-1}, \quad \mathcal{P}_{n-1,p+r(n-1,p)-\ell} \quad \text{and} \quad \mathcal{P}_{n-1,p+r(n-1,p-1)-\ell-1}. \qquad (8.23)$$

With this in hand, we can now attend to the proof of the theorem.

Proof of Theorem 8.2. Because of Proposition 8.3, we can (and will) assume throughout that $\ell > 2$. So recall that when $n = 1$, $\mathcal{P}_{1,0} = \mathsf{TL}_1$ is the only principal indecomposable and it coincides with $\mathcal{V}_{1,0}$ (and $\mathcal{L}_{1,0}$). This therefore agrees with the statement of the theorem. Clearly $\mathcal{L}_{1,0}$ is the unique irreducible, up to isomorphism.

The induction step over n needs to be supplemented by a further induction on the label of the critical strip k. The following statements are to be proved for all $k(n,p)$:

1) The submodule of $\mathcal{P}_{n-1,p}\!\uparrow$ which is the generalised eigenspace of F_n with eigenvalue $f_{n,p}$ is either $\mathcal{P}_{n,p}$ or, when (n,p) is critical, $\mathcal{P}_{n,p} \oplus \mathcal{P}_{n,p+\ell}$. In the latter case, $\mathcal{P}_{n,p} \cong \mathcal{V}_{n,p}$ and, in the former, $\mathcal{P}_{n,p}$ forms the exact sequence

$$\mathcal{V}_{n,p+r(n,p)} \xrightarrow{\iota} \mathcal{P}_{n,p} \longrightarrow \mathcal{V}_{n,p} \longrightarrow 0, \qquad (8.24)$$

where $\mathcal{V}_{n,p+r(n,p)}$ is $\{0\}$ when $(n,p+r(n,p))$ is outside the Bratteli diagram.
2) All copies of $\mathcal{P}_{n,p}$ in TL_n are induced from the copies of $\mathcal{P}_{n-1,p'}$ in TL_{n-1}, where

$$p' \in \begin{cases} \{p\} & \text{if } r(n,p) = \ell - 1, \\ \{p, p-1, p-\ell\} & \text{if } r(n,p) = \ell, \\ \{p, p-1\} & \text{otherwise.} \end{cases} \qquad (8.25)$$

To the left of, or on, the first critical line – We first consider the base case $k(n,p) = 0$ which labels all (n,p) to the left of and on the first critical line. We start with a pair $(n-1,p)$ lying strictly to the left of the first critical line on the Bratteli diagram, so $k(n-1,p) = 0$ and $r(n-1,p) < \ell$. We may assume, inductively, that $\mathcal{P}_{n-1,p} \cong \mathcal{V}_{n-1,p}$. Since $(n-1,p)$ is not critical, Corollary 6.6



Figure 3. The principal indecomposables that might contribute to $\mathcal{P}_{n,p}$ when $k(n,p) = 0$ for the four cases $r = r(n,p) = 1$, $1 < r < \ell - 1$, $r = \ell - 1$ and $r = \ell$. The thick vertical segments indicate the critical lines of the Bratteli diagrams, the dotted one the left boundary of the diagram. A curve between the symmetric pairs $(n-1, p'')$ and $(n-1, p')$, with $p'' > p'$, is drawn to remind us that, by the induction hypothesis, the projective module $\mathcal{P}_{n-1,p'}$ belongs to a non-split short exact sequence $0 \to V_{n-1,p''} \to \mathcal{P}_{n-1,p'} \to V_{n-1,p'} \to 0$.

gives $\mathcal{P}_{n-1,p}\!\uparrow\, \cong \mathcal{V}_{n,p+1} \oplus \mathcal{V}_{n,p}$ (with $\mathcal{V}_{n,\lfloor n/2 \rfloor+1} = \{0\}$). Thus, $\mathcal{V}_{n,p}$ is projective for $1 \leqslant r(n,p) \leqslant \ell$. Moreover, $\mathcal{V}_{n,p}$ is indecomposable, by Proposition 3.3, hence we conclude that $\mathcal{P}_{n,p} \cong \mathcal{V}_{n,p}$ for any p satisfying $k(n,p) = 0$. This proves (1).

We next determine, for $k(n,p) = 0$, which TL_{n-1} principal indecomposables contribute copies of $\mathcal{P}_{n,p} \cong \mathcal{V}_{n,p}$ to TL_n and which do not. The principal modules among those of (8.23) that can actually contribute to a given $\mathcal{P}_{n,p}$ depend on whether $r = r(n,p)$ is 1, strictly between 1 and $\ell - 1$, equal to $\ell - 1$ or ℓ. On each of the diagrams of Figure 3, the possible contributing projective modules are identified by their labels. For example, in the case $r = r(n,p) = 1$ which, when $k = 0$, corresponds to $(n,p) = (2p,p)$, only two of the four possibilities of (8.23) are distinct and non-vanishing: $\mathcal{P}_{2p-1,p-1} \cong \mathcal{V}_{2p-1,p-1}$ and $\mathcal{P}_{2p-1,p-\ell+1}$. Since we assume that $\ell > 2$, $(2p-1, p-1)$ is not critical, and the exact sequence for $\mathcal{V}_{2p-1,p-1}\!\uparrow$ splits. By Theorem B.3, there are $\dim \mathcal{L}_{2p-1,p-1}$ copies of $\mathcal{P}_{2p-1,p-1}$ in TL_{2p-1}, hence we obtain this many copies of $\mathcal{P}_{2p,p}$ in TL_{2p}. Corollary 5.2 now gives $\dim \mathcal{L}_{2p,p} = \dim \mathcal{L}_{2p-1,p-1}$, which is precisely the number of copies of $\mathcal{P}_{2p,p}$ that Theorem B.3 requires in TL_{2p}. This shows that every copy of $\mathcal{P}_{n,p}$ in TL_n, with $k(n,p) = 0$ and $r(n,p) = 1$, is induced from the copies of $\mathcal{P}_{n-1,p-1}$ in TL_{n-1} (and not from the copies of $\mathcal{P}_{n-1,p-\ell+1}$).

One similarly argues that every copy of $\mathcal{P}_{n,p}$ in TL_n, with $k(n,p) = 0$ and $1 < r(n,p) < \ell - 1$, is induced from the copies of $\mathcal{P}_{n-1,p}$ and $\mathcal{P}_{n-1,p-1}$ in TL_{n-1}. This essentially follows from $\dim \mathcal{L}_{n,p} = \dim \mathcal{L}_{n-1,p} + \dim \mathcal{L}_{n-1,p-1}$. When $k(n,p) = 0$ and $r(n,p) = \ell - 1$, one instead uses $\dim \mathcal{L}_{n,p} = \dim \mathcal{L}_{n-1,p}$ to conclude that every copy of $\mathcal{P}_{n,p}$ is induced from copies of $\mathcal{P}_{n-1,p}$. The



$r=1$ $\quad$ $1<r<\ell-1$

$r=\ell-1$ $\quad$ $r=\ell$

Figure 4. The principal indecomposables that might contribute to $\mathcal{P}_{n,p}$ when $k(n,p)>0$ and $r=r(n,p)=1$, $1<r<\ell-1$, $r=\ell-1$ and $r=\ell$. See Figure 3 for the notation.

calculation for $k(n,p)=0$ and $r(n,p)=\ell$ is a little more interesting. Then, (8.23) gives $\mathcal{P}_{n-1,p}\cong\mathcal{V}_{n-1,p}$, $\mathcal{P}_{n-1,p-1}$ and $\mathcal{P}_{n-1,p-\ell}$ as candidates for inducing $\mathcal{P}_{n,p}$. However, note that $\mathcal{P}_{n-1,p-1}$ could contribute *twice*, because the second exact sequence of (8.22) becomes $\mathcal{V}_{n,p}\to\mathcal{P}'\to\mathcal{V}_{n,p}\to 0$. In fact, the calculation

$$\begin{aligned}\dim\mathcal{L}_{n-1,p}+2\dim\mathcal{L}_{n-1,p-1}+\dim\mathcal{L}_{n-1,p-\ell}&=\dim\mathcal{L}_{n-1,p}+\dim\mathcal{R}_{n-1,p}+\dim\mathcal{L}_{n-1,p-1}+\dim\mathcal{R}_{n-1,p-1}\\&=\dim\mathcal{V}_{n-1,p}+\dim\mathcal{V}_{n-1,p-1}=\dim\mathcal{V}_{n,p}=\dim\mathcal{L}_{n,p}\end{aligned}\tag{8.26}$$

shows that $\mathcal{P}_{n-1,p-1}$ does contribute twice and $\mathcal{P}_{n-1,p}$ and $\mathcal{P}_{n-1,p-\ell}$ both contribute once each. Here, we have used Propositions 5.3 and 3.3, then Proposition 4.1 (or direct calculation using Equation (2.9)) and finally, Corollary 4.8. The above computation assumed implicitly that the n-th row of the Bratteli diagram is long enough to contain the labels $(n-1,p-1)$ and $(n-1,p-\ell)$. If it is not, then the following addition is necessary: If $(n,p)=(n,0)$, then only $\mathcal{P}_{n-1,p}$ contributes and it provides $\dim\mathcal{L}_{n-1,p}$ copies. But, in this case, $\dim\mathcal{L}_{n,0}=\dim\mathcal{L}_{n-1,0}$ and the result follows trivially. If (n,p) is such that $\mathcal{P}_{n-1,p-\ell}$ does not exist (because $p-\ell<0$), then the term $\dim\mathcal{R}_{n-1,p-1}$ in the first line of (8.26) should be removed. However it can be left there, as this radical is trivial and its dimension 0. This proves (2) and concludes the base case for the induction over $k(n,p)$.

To the right of the first critical line – We now suppose that $k(n,p)=k>0$, assuming that both (1) and (2) hold for all p' such that $k(n,p')<k$. Again, the diagrams of Figure 4 will be useful to identify the contributing modules.

If $r(n,p)=1$, then $(n-1,p)$ is critical, so $\mathcal{P}_{n-1,p}\!\uparrow\,\cong\mathcal{V}_{n-1,p}\!\uparrow$ has exact sequence $0\to\mathcal{V}_{n,p+1}\to\mathcal{P}_{n-1,p}\!\uparrow\,\to\mathcal{V}_{n,p}\to 0$. This cannot split, as otherwise $\mathcal{P}_{n,p+1}\cong\mathcal{V}_{n,p+1}$ and $\mathcal{P}_{n-1,p}$ would contribute a copy of $\mathcal{P}_{n,p+1}$, contradicting (2) (since $k(n,p+1)=k-1$ and $r(n,p+1)=\ell-1$). Thus,

$\mathcal{P}_{n,p} \cong \mathcal{P}_{n-1,p}\!\uparrow$, proving (1) for $k(n,p) = k$ and $r(n,p) = 1$. To prove (2), note that the above argument shows that $\mathcal{P}_{n-1,p}$ leads to $\dim \mathcal{L}_{n-1,p}$ copies of $\mathcal{P}_{n,p}$. To obtain further copies, we consider $\mathcal{P}_{n-1,p-1}$. This satisfies the exact sequence $0 \to \mathcal{V}_{n-1,p+1} \to \mathcal{P}_{n-1,p-1} \to \mathcal{V}_{n-1,p-1} \to 0$, from which we obtain the exactness of

$$\mathcal{V}_{n,p+1} \longrightarrow \mathcal{P}' \longrightarrow \mathcal{V}_{n,p} \longrightarrow 0 \tag{8.27}$$

as in (8.22). Since $\mathcal{P}'$ is projective with quotient $\mathcal{L}_{n,p}$, it is a direct sum of principal indecomposables, one of which must be $\mathcal{P}_{n,p}$. But, $\dim \mathcal{P}' \leqslant \dim \mathcal{V}_{n,p+1} + \dim \mathcal{V}_{n,p} = \dim \mathcal{P}_{n,p}$ because we have just established that $\mathcal{P}_{n,p} \cong \mathcal{P}_{n-1,p}\!\uparrow$. So we conclude that $\mathcal{P}' \cong \mathcal{P}_{n,p}$ and hence that $\mathcal{P}_{n-1,p-1}$ leads to $\dim \mathcal{L}_{n-1,p-1}$ more copies of $\mathcal{P}_{n,p}$. We therefore have all the required copies by Corollary 5.2 and Theorem B.3, proving (2) for $k(n,p) = k$ and $r(n,p) = 1$. (Note that, if $(n,p) = (n,0)$, the principal $\mathcal{P}_{n-1,p-1}$ is not defined. However, in this case, $\dim \mathcal{L}_{n,p} = \dim \mathcal{L}_{n-1,p}$ and the right number of copies had already been provided by $\mathcal{P}_{n-1,p}$.)

For $1 < r(n,p) < \ell - 1$, we start with the exact sequence $0 \to \mathcal{V}_{n-1,p+r(n-1,p)} \to \mathcal{P}_{n-1,p} \to \mathcal{V}_{n-1,p} \to 0$, obtaining the exactness of

$$\mathcal{V}_{n,p+r(n,p)} \xrightarrow{\iota} \mathcal{P} \longrightarrow \mathcal{V}_{n,p} \longrightarrow 0. \tag{8.28}$$

If this does not split, then combining Lemma 8.1 and Corollary 7.4 yields $\mathcal{P}_{n,p} \cong \mathcal{P}$ as required. If this does split, then $\mathcal{V}_{n,p+r(n,p)}/\ker \iota$ will be indecomposable and projective, hence can only be $\{0\}$ (so $\mathcal{P}_{n,p} \cong \mathcal{V}_{n,p}$ as (1) requires) or $\mathcal{P}_{n,p+r(n,p)}$. But, the latter possibility implies that $\mathcal{P}_{n-1,p}$ contributes copies of $\mathcal{P}_{n,p+r(n,p)}$, contradicting (2) (since $k(n,p+r(n,p)) = k-1$ and $r(n,p+r(n,p)) = \ell - r(n,p)$). This proves (1) when $k(n,p) = k$ and $1 < r(n,p) < \ell - 1$. Statement (2) is now proven by noting that this analysis provides $\dim \mathcal{L}_{n-1,p}$ copies of $\mathcal{P}_{n,p}$ and the same analysis applied to $\mathcal{P}_{n-1,p-1}$ provides the rest, by Corollary 5.2.

When $r(n,p) = \ell - 1$, we prove (1) in exactly the same manner as for $1 < r(n,p) < \ell - 1$, starting with $\mathcal{P}_{n-1,p}$. Proving (2) is even easier because Corollary 5.2 implies that in this case, $\mathcal{P}_{n-1,p}$ contributes all the copies of $\mathcal{P}_{n,p}$. We therefore turn to $r(n,p) = \ell$. The exactness of $0 \to \mathcal{V}_{n-1,p+\ell-1} \to \mathcal{P}_{n-1,p} \to \mathcal{V}_{n-1,p} \to 0$ now yields that of

$$\mathcal{V}_{n,p+\ell} \longrightarrow \mathcal{P} \longrightarrow \mathcal{V}_{n,p} \longrightarrow 0. \tag{8.29}$$

If $k = 1$, then $\mathcal{V}_{n,p+\ell}$ is not defined, hence $\mathcal{P}_{n,p} \cong \mathcal{V}_{n,p}$ as (1) requires. For $k > 1$, it is easily seen that $\mathcal{P}$ must split as $\mathcal{V}_{n,p+\ell} \oplus \mathcal{V}_{n,p}$ to avoid a contradiction with (2) — since $k(n,p+\ell) = k-2$ and $r(n,p+\ell) = \ell$, $\mathcal{P}_{n-1,p}$ is known to contribute copies of $\mathcal{P}_{n,p+\ell} \cong \mathcal{V}_{n,p+\ell}$ to TL_n. This proves (1) when $k(n,p) = k$ and $r(n,p) = \ell$, and (2) follows from the same argument used when $k(n,p) = 0$. This completes the (inductive) proof of statements (1) and (2).

Completing the short exact sequences – The first question to answer is whether the induction of the $\mathcal{P}_{n-1,p}$ allows us to construct *all* of the principal indecomposables of TL_n. By the previous recursive argument, all principal indecomposables of TL_n must be summands of one of the $\mathcal{P}_{n-1,p}\!\uparrow$. Again, each $\mathcal{P}_{n-1,p}\!\uparrow$ can be written as a direct sum of at most two generalized eigenspaces of F_n, each being projective and part of an exact sequence of the form

$$\mathcal{V}_{n,p''} \xrightarrow{\iota} \mathcal{P} \longrightarrow \mathcal{V}_{n,p'} \longrightarrow 0, \tag{8.30}$$

where $\mathcal{P}$ is either one of the two summands, or equivalently

$$0 \to \mathcal{V}_{n,p''}/\ker\iota \to \mathcal{P} \to \mathcal{V}_{n,p'} \to 0. \tag{8.31}$$

If this exact sequence does not split, then $\mathcal{P}$ is projective and indecomposable (by Lemma 8.1). It is therefore a principal indecomposable with irreducible quotient $\mathcal{L}_{n,p'}$ and must be identified with $\mathcal{P}_{n,p'}$ (which has already been constructed). If the sequence splits, then the two summands $\mathcal{V}_{n,p''}/\ker\iota$ and $\mathcal{V}_{n,p'}$ are likewise projective and indecomposable (by Proposition 3.3 or Corollary 7.4), so must again be principal indecomposables (if not trivial). In fact, they must be $\mathcal{P}_{n,p''}$ and $\mathcal{P}_{n,p'}$ since their irreducible quotients are $\mathcal{L}_{n,p''}$ and $\mathcal{L}_{n,p'}$ respectively. Hence the set of $\mathcal{P}_{n,p}$, $0 \leqslant p \leqslant \lfloor n/2 \rfloor$, is a complete set of non-isomorphic principal indecomposable modules of TL_n. It follows that the corresponding $\mathcal{L}_{n,p}$ also form a complete set of non-isomorphic irreducible TL_n-modules.

We have proved, with statement (1) for non-critical (n,p), that $\mathcal{P}_{n,p}$ is part of the short exact sequence

$$0 \to \mathcal{V}_{n,p+r(n,p)}/\ker\iota \to \mathcal{P}_{n,p} \to \mathcal{V}_{n,p} \to 0, \tag{8.32}$$

where ι is the map that makes the sequence (8.24) exact at $\mathcal{P}_{n,p}$. Therefore, the dimension of $\mathcal{P}_{n,p}$ is bounded by

$$\dim\mathcal{P}_{n,p} \leqslant \dim\mathcal{V}_{n,p} + \dim\mathcal{V}_{n,p+r(n,p)} \tag{8.33}$$

with the bound saturated if and only if $\ker\iota = \{0\}$. The last step of the proof is to prove that this bound is indeed reached and therefore that the short sequence (8.17) is exact.

Since the $\mathcal{P}_{n,p}$ form a complete set of principal indecomposables, their dimensions are constrained by Theorem B.3:

$$\dim\mathsf{TL}_n = \sum_{p=0}^{\lfloor n/2 \rfloor} (\dim\mathcal{V}_{n,p})^2 = \sum_{p=0}^{\lfloor n/2 \rfloor} \dim\mathcal{L}_{n,p}\,\dim\mathcal{P}_{n,p}. \tag{8.34}$$

Some of the terms in these two sums coincide pairwise. Indeed, when (n,p) is critical, $\dim\mathcal{V}_{n,p} = \dim\mathcal{L}_{n,p} = \dim\mathcal{P}_{n,p}$. Let (n,p_1) be on the left of the first critical line of the Bratteli diagram and let $p_1 > p_2 > \cdots > p_m \geq 0$ be all the p_i obtained from p_1 by reflections with respect to critical lines. There is precisely one p_i between each pair of consecutive critical lines. The set $\{(n,p_i), 1 \leqslant i \leqslant m\}$ is the orbit of (n,p_1) under these reflections and every non-critical (n,p) belongs to precisely one such orbit.[8] Theorem 7.2 and the definition of $\mathcal{L}_{n,p}$ give

$$\dim\mathcal{L}_{n,p_{i+1}} = \dim\mathcal{R}_{n,p_i} \qquad \text{and} \qquad \dim\mathcal{L}_{n,p_1} = \dim\mathcal{V}_{n,p_1} - \dim\mathcal{R}_{n,p_1}. \tag{8.35}$$

Since the radical $\mathcal{R}_{n,p}$ is trivial for all p that do not have a reflection through a critical line on their right, we also have

$$\dim\mathcal{V}_{n,p_m} = \dim\mathcal{L}_{n,p_m}. \tag{8.36}$$

---

[8] These orbits were introduced in [12] and play a central role in the construction of the blocks of TL_n.

These identities, together with the bound (8.33), lead to the following bound for the right sum of (8.34) restricted to the orbit of (n,p_1):

$$\begin{aligned}\sum_{i=1}^{m}\dim\mathcal{L}_{n,p_i}\,\dim\mathcal{P}_{n,p_i} &= \dim\mathcal{L}_{n,p_1}\dim\mathcal{P}_{n,p_1}+\sum_{i=2}^{m}\dim\mathcal{L}_{n,p_i}\dim\mathcal{P}_{n,p_i}\\ &\leqslant (\dim\mathcal{V}_{n,p_1}-\dim\mathcal{R}_{n,p_1})\dim\mathcal{V}_{n,p_1}+\sum_{i=2}^{m}\dim\mathcal{L}_{n,p_i}(\dim\mathcal{V}_{n,p_i}+\dim\mathcal{V}_{n,p_{i-1}})\\ &= (\dim\mathcal{V}_{n,p_1})^2-\dim\mathcal{R}_{n,p_1}\dim\mathcal{V}_{n,p_1}+\sum_{i=2}^{m}\dim\mathcal{L}_{n,p_i}\dim\mathcal{V}_{n,p_i}+\sum_{i=1}^{m-1}\dim\mathcal{R}_{n,p_i}\dim\mathcal{V}_{n,p_i}\\ &= (\dim\mathcal{V}_{n,p_1})^2+\sum_{i=2}^{m-1}(\dim\mathcal{L}_{n,p_i}+\dim\mathcal{R}_{n,p_i})\dim\mathcal{V}_{n,p_i}+\dim\mathcal{L}_{n,p_m}\dim\mathcal{V}_{n,p_m}\\ &= \sum_{i=1}^{m}(\dim\mathcal{V}_{n,p_i})^2. \qquad (8.37)\end{aligned}$$

For (8.34) to hold, the upper bound must be reached for all orbits and the short sequence (8.17) must thus be exact for all $\mathcal{P}_{n,p}$ to the right of the first critical line. ∎

9 Summary of Results

We conclude the article with a brief summary outlining what has been proven and a few ideas concerning what can be done with the results. First, we have presented the well-known equivalence between the algebraic and diagrammatic definitions of the Temperley-Lieb algebra TL_n (with parameter $\beta = q + q^{-1}$), deducing as a consequence that its dimension is given by the n-th Catalan number. We have then discussed the standard TL_n-modules $\mathcal{V}_{n,p}$, with $0 \leqslant p \leqslant \lfloor n/2 \rfloor$, each of which admits a natural invariant bilinear form, and explained that a study of the irreducibility of the $\mathcal{V}_{n,p}$ may be reduced, in almost all cases, to the consideration of the non-degeneracy of this bilinear form. Of course, we have also detailed how to analyse the exceptional cases when the natural bilinear form is not useful.

The question of whether these standard modules are irreducible or not turned out to have structural implications for the Temperley-Lieb algebra. We used restriction from TL_n to TL_{n-1} to derive a recursion relation describing the kernel of the bilinear form on each $\mathcal{V}_{n,p}$. This gave a complete answer to the irreducibility of the $\mathcal{V}_{n,p}$ and led to a criterion for the semisimplicity of TL_n. In particular, this recovered the well-known result that the TL_n are all semisimple when q is not a root of unity. While our strategy follows that of [13] rather closely, we believe that our proof is new. Moreover, we found that the aforementioned description of the kernel suggested that the standard modules were either irreducible or had an irreducible maximal submodule, the radical.

To prove this suggestion, we turned to the induced modules obtained from the $\mathcal{V}_{n,p}$ by including TL_n in TL_{n+1} in the obvious way. After deriving the required structural information concerning these induced modules, we detailed a non-trivial computation involving the action of a little-known central element $F_n \in \mathsf{TL}_n$ on the induced modules. This then allowed us to deduce the existence of non-trivial homomorphisms between certain standard modules. As a consequence, we obtained the desired irreducibility of the (non-trivial) radicals as well as a complete description of the space of homomorphisms between any two standard modules. All of these proofs appear to be new.

Finally, we used our knowledge of induced modules to systematically construct the principal indecomposable modules $\mathcal{P}_{n,p}$ for the Temperley-Lieb algebra. These turn out to be realised as either standard modules or as submodules, corresponding to a given generalised eigenspace of the central element F_n, of a multiply-induced standard module. The structures of the $\mathcal{P}_{n,p}$ followed immediately from the analysis, as did the statement that the irreducible TL_n-modules $\mathcal{L}_{n,p}$ that can be constructed from the $\mathcal{V}_{n,p}$ are, in fact, exhaustive (up to isomorphism). While the structure of the principal indecomposables is known (it appeared first in [2]), we are confident that our straightforward proof is also new. Certainly, it relies upon a particularly remarkable property of F_n, namely that it completely distinguishes the non-critical blocks of TL_n, a property which does not seem to be shared by other better-known central elements.

We include, for convenience, pictorial representations of the structures of the standard and principal indecomposable modules when q is a root of unity. These take the form of annotated Loewy diagrams which are popular in, for example, logarithmic conformal field theory. The Loewy diagrams for the standard modules have two forms according as to whether (n,p) is *critical*, meaning $q^{2(n-2p+1)} = 1$, or not. Let (n,p) and (n,p') form a symmetric pair, in the sense of Section 7, with $p > p'$. The possible Loewy diagrams for the $\mathcal{V}_{n,p}$ $(0 \leqslant p \leqslant \lfloor n/2 \rfloor)$ are then as follows:

$$\mathcal{L}_{n,p}$$

$\mathcal{V}_{n,p}$

(n,p) critical

$$\begin{tikzcd} \mathcal{L}_{n,p} \arrow[d] \\ \mathcal{L}_{n,p'} \end{tikzcd}$$

$\mathcal{V}_{n,p}$

(n,p) non-critical
$(n \neq 2p$ or $\beta \neq 0)$

$$\mathcal{L}_{2p,p-1}$$

$\mathcal{V}_{n,p}$

$(n = 2p$ and $\beta = 0)$.

To explain, these diagrams present the (irreducible) composition factors of the $\mathcal{V}_{n,p}$ as nodes, connected by arrows, that are meant to represent the action of the algebra. The diagrams on the left and right indicate that $\mathcal{V}_{n,p}$ is in fact irreducible (and isomorphic to $\mathcal{L}_{n,p}$ and $\mathcal{L}_{2p,p-1}$, respectively) in these cases. By contrast, the diagram in the middle has its only arrow pointing towards $\mathcal{L}_{n,p'}$, indicating that $\mathcal{V}_{n,p}$ has, in this case, a submodule isomorphic to $\mathcal{L}_{n,p'}$. In fact, it is the maximal submodule — quotienting corresponds to removing the node $\mathcal{L}_{n,p'}$ and the arrow, leaving us with only $\mathcal{L}_{n,p}$ (and no arrows). In other words, $\mathcal{V}_{n,p}/\mathcal{L}_{n,p'} \cong \mathcal{L}_{n,p}$. We remark that for sufficiently small p, the p' required to form a symmetric pair would be negative. In that case, $\mathcal{L}_{n,p'}$ should be understood to be $\{0\}$. The diagram in the middle should then be taken to degenerate into the diagram on the left.

The Loewy diagrams for the principal indecomposables are only a little more complicated. First, we remark that when $\beta = 0$ and n is even, p is restricted to the range $0 \leqslant p \leqslant n/2 - 1$; otherwise, the range is $0 \leqslant p \leqslant \lfloor n/2 \rfloor$. Now, take $p'' > p > p'$ so that (n,p'') and (n,p), as well as (n,p) and (n,p'), form symmetric pairs. The possible Loewy diagrams for the $\mathcal{P}_{n,p}$ are then as follows:

$$\begin{tikzcd}
\mathcal{L}_{n,p}
\end{tikzcd}$$

$\mathcal{P}_{n,p}$

(n,p) critical

$$\begin{tikzcd}
& \mathcal{L}_{n,p} \arrow[dl] \arrow[dr] & \\
\mathcal{L}_{n,p''} \arrow[dr] & & \mathcal{L}_{n,p'} \arrow[dl] \\
& \mathcal{L}_{n,p} &
\end{tikzcd}$$

$\mathcal{P}_{n,p}$

(n,p) non-critical
$(n \neq 2p+2$ or $\beta \neq 0)$

$$\begin{tikzcd}
\mathcal{L}_{2p+2,p} \arrow[dr] & \\
& \mathcal{L}_{2p+2,p-1} \arrow[dl] \\
\mathcal{L}_{2p+2,p} &
\end{tikzcd}$$

$\mathcal{P}_{n,p}$

$(n = 2p+2$ and $\beta = 0)$.

Again, the diagram on the left indicates that the critical principal indecomposables are irreducible. To understand the diagram in the middle, note that there is a submodule isomorphic to $\mathcal{L}_{n,p}$ (the bottom node with arrows only pointing in). Quotienting by this submodule leads to a module with three composition factors. It has a submodule isomorphic to $\mathcal{L}_{n,p''} \oplus \mathcal{L}_{n,p'}$,[9] and quotienting by this submodule results in something isomorphic to $\mathcal{L}_{n,p}$. The diagram on the right is interpreted similarly.

The degeneracy conditions are likewise a little more complicated: When p is small enough in the middle diagram that p' would have to be negative, one must delete the node $\mathcal{L}_{n,p'}$ and arrows pointing towards or away from it. The resulting principal indecomposable therefore has three composition factors. However, when p is large enough that p'' would have to be greater than $\lfloor n/2 \rfloor$, one should delete *both* $\mathcal{L}_{n,p''}$ and the bottom $\mathcal{L}_{n,p}$, so that $\mathcal{P}_{n,p}$ has only two composition factors. This means that $\mathcal{P}_{n,p} \cong \mathcal{V}_{n,p}$ when p is on the left of the first critical line. It may, of course, happen for n sufficiently small that there are non-critical p for which both p' and p'' fall outside the allowed range. Then, the middle diagram degenerates into the diagram on the left. Similarly, the diagram on the right degenerates only when $n = 2$ and $p = 0$. In this case, the Loewy diagram for $\mathcal{P}_{2,0}$ has two composition factors, both isomorphic to $\mathcal{L}_{2,0}$, connected by an arrow.

---

[9] One can (and should) ask why this submodule is a direct sum. Equivalently, why is there no arrow from the factor $\mathcal{L}_{n,p'}$ to the factor $\mathcal{L}_{n,p''}$ in the middle diagram? (That there is no arrow in the opposite direction follows directly from (8.17).) This may be answered by showing that any short exact sequence of the form $0 \to \mathcal{L}_{n,p''} \to \mathcal{M} \to \mathcal{L}_{n,p'} \to 0$ necessarily splits. To see this, note that projectivity implies the existence of a homomorphism $\delta \colon \mathcal{P}_{n,p'} \to \mathcal{M}$ such that the following diagram commutes:

$$\begin{tikzcd}
& & \mathcal{P}_{n,p'} \arrow[d, "\delta"'] \arrow[dr] & & \\
0 \arrow[r] & \mathcal{L}_{n,p''} \arrow[r] & \mathcal{M} \arrow[r] & \mathcal{L}_{n,p'} \arrow[r] & 0.
\end{tikzcd}$$

As $\mathcal{M}$ has two composition factors, there are four possibilities for $\mathrm{im}(\delta)$ and it is easily checked that only $\mathrm{im}(\delta) \cong \mathcal{L}_{n,p'}$ is consistent. Thus, $\mathcal{L}_{n,p'}$ is a submodule of $\mathcal{M}$, so $\mathcal{M} \cong \mathcal{L}_{n,p''} \oplus \mathcal{L}_{n,p'}$.

Finally, we remark that such a complete description of the principal indecomposables allows one to apply standard tools from homological algebra to answer more advanced questions regarding the variety of possible indecomposable structures. We shall not do so here, but will content ourselves with mentioning that it is now easy to write down projective presentations and resolutions for standard and irreducible modules and so compute the extension groups between them. This technology answers, in particular, the question of whether giving a non-split short exact sequence completely determines the module in the middle. One can also use the extension computations to construct injective modules and, indeed, give a complete classification of the indecomposable modules of the Temperley-Lieb algebra (something which is not possible for most associative algebras). These directions are clearly crucial for a sound mathematical understanding of Temperley-Lieb representation theory as well as necessary for a complete identification of Temperley-Lieb modules in physical models. In both cases, the structural results proven here provide a springboard from which one can profitably tackle advanced questions.

Acknowledgements

The authors would like to thank Alexi Morin-Duchesne for a useful suggestion and Azat Gainutdinov, Gus Lehrer and Paul Martin for helpful discussions. DR is supported by an Australian Research Council Discovery Project DP0193910 and YSA holds a grant from the Canadian Natural Sciences and Engineering Research Council. This support is gratefully acknowledged.

Appendix A Central Elements

The goal of this appendix is to introduce the central element $F_n \in \mathsf{TL}_n$ which plays a crucial role in Sections 7 and 8. This is not the only important central element in the theory of the Temperley-Lieb algebra. One should also mention the well-known central element $C_n = (t_1 \cdots t_{n-1})^n \in \mathsf{TL}_n$, where $t_i = \mathbf{1} - qu_i$, which is derived from its analogue in the braid group [22, 23], as well as the central idempotents described by Jones [5] and Wenzl [11]. For our purposes, F_n will suffice (whereas C_n in particular will not). This appendix uses notation and one result from each of Sections 2 and 3; results from other sections are also used to provide examples. The element F_n is first used in Corollary 4.2.

The role of central elements is well-known to physicists, where the quadratic Casimir of a semisimple Lie algebra probably provides the most familiar example. One use for such central elements is to decompose representations into their eigenspaces. More precisely, if c is a central element of an algebra A and $\mathcal{M}$ a module over this algebra, then the linear map $\phi_c \colon \mathcal{M} \to \mathcal{M}$ obtained by left-multiplying elements of $\mathcal{M}$ by c defines a homomorphism of modules: $a\phi_c(m) = acm = cam = \phi_c(am)$ for all $a \in \mathsf{A}$, $m \in \mathcal{M}$. The eigenspaces of ϕ_c are then submodules of $\mathcal{M}$. However, it might happen that ϕ_c is not diagonalisable on $\mathcal{M}$, in which case the generalised eigenspaces $\mathcal{M}_\lambda = \{m \in \mathcal{M} \mid (\phi_c - \lambda\mathbf{1})^{\dim \mathcal{M}} m = 0\}$ are also submodules. In fact, $\mathcal{M}$ splits, as an A-module, into the *direct sum* of the generalised eigenspaces of ϕ_c. This is a very useful property.

We therefore turn to the central element $F_n \in \mathsf{TL}_n$ and its properties. The computations which follow, and even the definition of F_n itself, are diagrammatic. The construction is based upon the recent analysis of a double row transfer matrix [8]. It also appears briefly in [24] (with $\beta = 0$), though it does not seem to have been recognised as central there. This important property was made explicit in [25], where its eigenvalues were computed. We will follow their approach with only minor changes.

To define F_n in TL_n, it is useful to introduce the following shorthand (recall that $\beta = q + q^{-1}$):



(A.1)

These *crossings* are to be interpreted as formal objects which will be used as building blocks to form (linear combinations of) n-diagrams. In particular, we define F_n as follows:

$F_n =$



(A.2)

This looks deceptively simple, but the notation hides a large number of diagrams as each crossing of thin lines stands for a sum of two terms. Computing even F_3 explicitly requires considering a linear combination of $2^6 = 64$ 3-diagrams, even though $\dim \mathsf{TL}_3 = 5$. For comparison, we list the first few explicit forms for F_n and the better known C_n:

$$F_1 = (q^2+q^{-2})\mathbf{1}, \qquad C_1 = \mathbf{1}, \tag{A.3a}$$

$$F_2 = (q^3+q^{-3})\mathbf{1} - (q-q^{-1})^2 u_1, \qquad C_2 = \mathbf{1} + q^2(q-q^{-1})u_1, \tag{A.3b}$$

$$\begin{aligned} F_3 &= (q^4+q^{-4})\mathbf{1} - (q-q^{-1})(q^2-q^{-2})(u_1+u_2) & C_3 &= \mathbf{1} + q^3(q^2-q^{-2})(u_1+u_2) \\ &\quad + (q-q^{-1})^2(u_1u_2+u_2u_1) & &\quad - q^3(q-q^{-1})(u_1u_2+u_2u_1). \end{aligned} \tag{A.3c}$$

We note that because the number of crossings used to construct F_n is even, the coefficients of the corresponding linear combination of n-diagrams will only contain integral powers of q and q^{-1}. Note further that



(A.4)

recalling that the adjoint was chosen in Section 3 to be linear, not antilinear. It follows that F_n is *self-adjoint*. Finally, it is also true [25] that F_n is invariant under $q \leftrightarrow q^{-1}$ (we remark that C_n does not have this last property).

Proposition A.1. *$F_n \in \mathsf{TL}_n$ is central.*

Proof. The key insight is that



(A.5a)

and similarly, that



(A.5b)

Thus,



(A.6)

is symmetric under a vertical reflection, hence self-adjoint. It follows that u_iF_n must be self-adjoint for all i, whence

$$u_iF_n = (u_iF_n)^\dagger = F_n^\dagger u_i^\dagger = F_nu_i, \tag{A.7}$$

as required. ■

The eigenvalue of C_n on the standard module $\mathcal{V}_{n,p}$ is easily shown to be $q^{2p(n+1-p)}$ (see for example [13]). The corresponding result for F_n is as follows.

Proposition A.2. *The element F_n acts on $\mathcal{V}_{n,p}$ as the identity times $f_{n,p} = q^{n-2p+1} + q^{-(n-2p+1)}$.*

Proof. First note that any central element must act as a multiple of the identity on $\mathcal{V}_{n,p}$ by Proposition 3.8. We may therefore determine this multiple by computing the action of F_n on any (n,p)-link state (modulo terms which have more than p links). A convenient choice is the state $z_p \in \mathcal{V}_{n,p}$ which has first p simple links, followed by $n-2p$ defects.

We will break the computation up into two pieces, corresponding to the action on the p simple links and the action on the $n-2p$ defects. The first is easy: Using Equation (A.5) (or rather its adjoint) repeatedly, we obtain



(A.8)

This will clearly contribute 1 or β, according as to whether the loop wiggling down is closed by the action on the defects.

This latter action requires a bit of explanation. First, apply the definition (A.1) to expand the top-right partial crossing (the loop at the top of these diagrams stands for the big loop of (A.8)):



(A.9)

Consider now the first term on the right-hand side. When we expand the crossing immediately below that just considered, only one term contributes. The other leads to a link on the middle vertical border which creates an extra link in the result (using the adjoint of Equation (A.5a)), so the result vanishes. This observation propagates down the right side, yielding a factor of $q^{1/2}$ at each step. Thus,

$q^{1/2}$  $= q^{(n-2p)/2}$  . (A.10)

Now expand the crossing at the top-left. Because of the big loop at the top, we again find that only one term contributes — the other corresponds to closing two defects and hence vanishes. This observation propagates down the left side, picking up a factor of $q^{1/2}$ at each step, until we reach the last crossing for which *both* terms contribute:

$q^{(n-2p)/2}$  $= q^{n-2p-1/2}$  $= q^{n-2p}$  $- q^{n-2p-1}$  . (A.11)

The overall contribution from this analysis is therefore a factor of $q^{n-2p}\beta - q^{n-2p-1} = q^{n-2p+1}$.

The analysis of the second term on the right-hand side of (A.9) is almost identical. After analysing the crossings on the right side, the diagrams become horizontal reflections of those we have just analysed (this reflection corresponds to the automorphism $u_i \leftrightarrow u_{n-i}$) and q is replaced by q^{-1}. The resulting contribution is therefore $q^{-(n-2p+1)}$. Summing the contributions from both terms now gives the desired result. ∎

We conclude by indicating why we consider the central element F_n superior to the more familiar C_n for the purpose of analysing the indecomposable structure of TL_n-modules. The algebra TL_n is semisimple when q is not a root of unity by Corollary 4.6, so we may as well suppose that there exists a minimal positive integer ℓ such that $q^{2\ell} = 1$. In other words, we have $q = e^{\mathrm{i}\pi k/\ell}$ for some $k \in \mathbb{Z}$ coprime to ℓ. The eigenvalue of F_n on the standard module $\mathcal{V}_{n,p}$ is therefore

$$f_{n,p} = 2\cos\frac{\pi k\,(n-2p+1)}{\ell}. \tag{A.12}$$

In Section 4, we defined (n,p) to be critical when ℓ divides $n-2p+1$. In that case, $f_{n,p}$ achieves it maximal or minimal value ± 2. More generally, we defined critical lines that partition the Bratteli diagram of the Temperley-Lieb algebras in Section 5. The crucial property of F_n is that if (n,p) and (n,p') are distinct pairs, both lying in the same strip bounded by two consecutive critical lines, then $f_{n,p} \neq f_{n,p'}$. In other words, the eigenvalue of F_n completely distinguishes standard modules within such strips. In the language of Goodman and Wenzl [12], this means that F_n completely characterises the non-critical *blocks* of the Temperley-Lieb algebra TL_n. This property is not shared by C_n. A simple example is provided at $q = e^{\mathrm{i}\pi/6}$ for TL_4. The three modules $\mathcal{V}_{4,2}$, $\mathcal{V}_{4,1}$ and $\mathcal{V}_{4,0}$

lie to the left of the first critical line. The eigenvalues of C_4 on each are 1, $e^{4\pi \mathrm{i}/3}$ and 1, and fail to distinguish $\mathcal{V}_{4,2}$ and $\mathcal{V}_{4,0}$. The eigenvalues of F_4 are $-\sqrt{3}$, 0 and $\sqrt{3}$.

Appendix B Basic results

This Appendix reviews certain classical tools of representation theory that are not commonly seen in a first course on representation theory, say at the level of [26]. These are Wedderburn's theorem and its generalisation to non-semisimple algebras, Frobenius reciprocity, the right-exactness of induction and the basic properties of projective modules. We state them here without proof. The reader can find proofs of Wedderburn's theorem in [27, 28]. Proofs of Frobenius reciprocity for algebras may be found in [2, 28]. The right-exactness of tensor products is proven in [29] and projective modules are covered in [21,28,30]. Finally, we note that [6, App. A] provides a concise introduction to finite-dimensional algebras that gives proofs of Wedderburn's theorem and its non-semisimple generalisation as well as what is needed on projective modules in order to understand Section 8. All modules in this summary should be understood to be finite-dimensional left modules.

Let A be a finite-dimensional associative algebra over $\mathbb{C}$. Every element of A can be seen to act on A by left multiplication. This action makes A into a left A-module called the *regular module*. The algebra A is said to be semisimple if the regular module is completely reducible, that is, if it can be written as a direct sum of irreducible modules. (In the case of a finite group G, its group algebra $\mathbb{C}G$ is always semisimple [26].) A key consequence of semisimplicity is the following theorem.

Theorem B.1 (Wedderburn)**.** *Let A be a complex, finite-dimensional, semisimple, associative algebra. Then, the regular module decomposes as*

$$\mathsf{A} \cong \bigoplus_{i=1}^{r} (\dim \mathcal{L}_i)\, \mathcal{L}_i, \tag{B.1}$$

where the $\mathcal{L}_1, \dots, \mathcal{L}_r$ form a complete set of non-isomorphic irreducible A-modules. That is, A decomposes as the direct sum of irreducibles with each irreducible appearing with multiplicity equal to its dimension.

A useful "converse" to this result which will be invoked in Section 4 is the following:

Proposition B.2. *If the regular representation of a complex, finite-dimensional, associative algebra A decomposes as in* (B.1)*, where the $\mathcal{L}_1, \dots, \mathcal{L}_r$ form a complete set of non-isomorphic irreducible A-modules, then A is semisimple.*

When the algebra A is not semisimple, the decomposition of its regular module will contain modules that are reducible, but not completely reducible. An *indecomposable* module $\mathcal{M}$ is one which cannot be written as the direct sum of two proper non-trivial submodules. Irreducible modules are examples. When the regular module of an algebra is written as a direct sum of indecomposable modules, those that appear in the decomposition are called the *principal indecomposable modules* of the algebra. Let $\{\mathcal{P}_i\}$ and $\{\mathcal{L}_i\}$ be complete sets of non-isomorphic principal indecomposable modules of A and non-isomorphic irreducible modules, respectively. Wedderburn's theorem does not hold when semisimplicity is relaxed. It is replaced by the following generalisation:

Theorem B.3. *Let* A *be a complex, finite-dimensional, associative algebra. The two sets* $\{\mathcal{P}_i\}$ *and* $\{\mathcal{L}_i\}$ *are put in one-to-one correspondence by associating a given principal indecomposable with its unique irreducible quotient. If* r *is the common cardinality of these sets, then the regular representation decomposes as*

$$\mathsf{A} \cong \bigoplus_{i=1}^{r} (\dim \mathcal{L}_i)\, \mathcal{P}_i. \tag{B.2}$$

The theorem of reciprocity due to Frobenius is the second result covered in this appendix. It relates modules constructed from known ones through the classical constructions of *restriction* and *induction*. The first is easy to describe: When the algebra A contains a subalgebra B, any module $\mathcal{M}$ over A is also a module over B. (The action of an element of B on $\mathcal{M}$ is simply that of this element seen as an element of A.) We will denote this B-module by $\mathcal{M}{\downarrow}$. As vector spaces, $\mathcal{M}$ and $\mathcal{M}{\downarrow}$ are identical and have the same dimension. The second construction, induction, also uses a pair of algebras $\mathsf{B} \subset \mathsf{A}$ as before, but the starting module $\mathcal{M}$ is now over B. The induced A-module, which we denote by $\mathcal{M}{\uparrow}$, is the tensor product $\mathsf{A} \otimes_{\mathsf{B}} \mathcal{M}$. (Section 6 provides a more detailed definition and several explicit examples.) Restriction and induction are "dual" operations in the following sense.

Proposition B.4 (Frobenius reciprocity)**.** *Let* $\mathsf{B} \subset \mathsf{A}$ *be two finite-dimensional associative algebras over* $\mathbb{C}$*. Let* $\mathcal{M}$ *be a* B*-module and* $\mathcal{N}$ *be an* A*-module. Then, the following isomorphism between vector spaces of module homomorphisms holds:*

$$\mathrm{Hom}_{\mathsf{A}}(\mathcal{M}{\uparrow}, \mathcal{N}) \cong \mathrm{Hom}_{\mathsf{B}}(\mathcal{M}, \mathcal{N}{\downarrow}).$$

The most familiar version of reciprocity corresponds to taking B and A to be the group algebras $\mathbb{C}H$ and $\mathbb{C}G$ of a pair of finite groups $H \subset G$. Finally, we shall need the behaviour of induction with respect to exact sequences.

Proposition B.5. *Suppose that*

$$0 \longrightarrow \mathcal{M}_1 \longrightarrow \mathcal{M}_2 \longrightarrow \mathcal{M}_3 \longrightarrow 0$$

is an exact short sequence of B*-modules. If* $\mathsf{B} \subset \mathsf{A}$*, then there is an exact sequence involving the induced* A*-modules* $\mathcal{M}_i{\uparrow} = \mathsf{A} \otimes_{\mathsf{B}} \mathcal{M}_i$*:*

$$\mathcal{M}_1{\uparrow} \longrightarrow \mathcal{M}_2{\uparrow} \longrightarrow \mathcal{M}_3{\uparrow} \longrightarrow 0.$$

Because induction preserves the exactness of short exact sequences except at the leftmost position, induction is said to be right-exact.

We turn now to projective modules which play a central role in Section 8. An A-module $\mathcal{P}$ is projective if, when there are two other A-modules $\mathcal{M}$ and $\mathcal{N}$ and homomorphisms $\alpha : \mathcal{P} \to \mathcal{N}$ and

$\gamma : \mathcal{M} \to \mathcal{N}$ with γ surjective, then there exists a homomorphism $\delta : \mathcal{P} \to \mathcal{M}$ such that $\gamma \circ \delta = \alpha$. Equivalently, with the same input, there exists a homomorphism $\delta : \mathcal{P} \to \mathcal{M}$ such that the following diagram, with the bottom row exact, commutes:

$$\begin{tikzcd}
& \mathcal{P} \arrow[dl, dashed, "\delta"'] \arrow[d, "\alpha"] & \\
\mathcal{M} \arrow[r, "\gamma"'] & \mathcal{N} \arrow[r] & 0.
\end{tikzcd}$$

Here are some basic properties of projective modules.

Proposition B.6.

(i) Any direct sum of projective modules is projective.
(ii) Every direct summand of a projective module is projective.

Clearly *(ii)* implies that every projective module can be written as a sum of indecomposable projective modules. There are many other ways to characterise projective modules.

Proposition B.7. *Let $\mathcal{P}$ be a module over an associative algebra. The three following conditions are equivalent:*

(i) $\mathcal{P}$ is projective.
(ii) if $\mathcal{M}$ and $\mathcal{N}$ are any modules such that the sequence $0 \to \mathcal{M} \to \mathcal{N} \to \mathcal{P} \to 0$ is exact, then this sequence splits.
(iii) $\mathcal{P}$ is a direct sum of principal indecomposable modules.

The last statement shows the particular role played by the summands of (B.2): The principal indecomposables are precisely the indecomposable projective modules.

References

[1] H Temperley and E Lieb. Relations Between the ‘Percolation’ and ‘Colouring’ Problem and Other Graph-Theoretic Problems Associated with Regular Plane Lattices: Some Exact Results for the ‘Percolation’ Problem. *Proc. Roy. Soc. London Ser. A*, 322:251–280, 1971.

[2] P Martin. *Potts Models and Related Problems in Statistical Mechanics*, volume 5 of *Advances in Statistical Mechanics*. World Scientific, Singapore, 1991.

[3] R Baxter. *Exactly Solved Models in Statistical Mechanics.* Academic Press, London, 1982.

[4] G Andrews, R Baxter, and P Forrester. Eight-Vertex SOS Model and Generalized Rogers-Ramanujan-Type Identities. *J. Stat. Phys.*, 35:193–266, 1984.

[5] V Jones. Index for Subfactors. *Invent. Math.*, 72:1–25, 1983.

[6] A Mathas. *Iwahori-Hecke Algebras and Schur Algebras*, volume 15 of *University Lecture Series*. AMS, Providence, 1998.

[7] A Gainutdinov, N Read, and H Saleur. Continuum Limit and Symmetries of the Periodic $gl(1|1)$ Spin Chain. arXiv:1112.3403 [hep-th].

[8] P Pearce, J Rasmussen, and J-B Zuber. Logarithmic Minimal Models. *J. Stat. Mech.*, 0611:P11017, 2006. arXiv:hep-th/0607232.

[9] N Read and H Saleur. Associative-Algebraic Approach to Logarithmic Conformal Field Theories. *Nucl. Phys.*, B777:316–351, 2007. arXiv:hep-th/0701117.

[10] P Mathieu and D Ridout. From Percolation to Logarithmic Conformal Field Theory. *Phys. Lett.*, B657:120–129, 2007. arXiv:0708.0802 [hep-th].

[11] H Wenzl. Hecke Algebras of type A_n and Subfactors. *Invent. Math.*, 92:249–383, 1988.

[12] F Goodman and H Wenzl. The Temperley-Lieb Algebra at Roots of Unity. *Pac. J. Math.*, 161:307–334, 1993.

[13] B Westbury. The Representation Theory of the Temperley-Lieb Algebras. *Math. Zeit.*, 219:539–565, 1995.

[14] G James and G Murphy. The Determinant of the Gram Matrix for a Specht Module. *J. Alg.*, 59:222–235, 1979.

[15] J Cardy. SLE for Theoretical Physicists. *Ann. Phys.*, 318:81–118, 2005. arXiv:cond-mat/0503313.

[16] A Gainutdinov and R Vasseur. Lattice Fusion Rules and Logarithmic Operator Product Expansions. arXiv:1203.6289 [hep-th].

[17] J Graham and G Lehrer. The Representation Theory of Affine Temperley-Lieb Algebras. *Enseign. Math.*, 44:173–218, 1998.

[18] P Di Francesco, O Golinelli, and E Guitter. Meanders and the Temperley-Lieb Algebra. *Comm. Math. Phys.*, 186:1–59, 1997. arXiv:hep-th/9602025.

[19] J Graham and G Lehrer. Cellular Algebras. *Invent. Math.*, 123:1–34, 1996.

[20] K Kytölä and D Ridout. On Staggered Indecomposable Virasoro Modules. *J. Math. Phys.*, 50:123503, 2009. arXiv:0905.0108 [math-ph].

[21] C Curtis and I Reiner. *Representation Theory of Finite Groups and Associative Algebras*, volume 11 of *Pure and Applied Mathematics*. Wiley, New York, 1962.

[22] W Chow. On the Algebraic Braid Group. *Ann. Math.*, 49:654–658, 1948.

[23] J Birman. *Braids, Links, and Mapping Class Groups*, volume 82 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, 1974.

[24] P Pearce, J Rasmussen, and S Villani. Solvable Critical Dense Polymers on the Cylinder. *J. Stat. Mech.*, 1002:P02010, 2010. arXiv:0910.4444 [hep-th].

[25] A Morin-Duchesne and Y Saint-Aubin. The Jordan Structure of Two Dimensional Loop Models. *J. Stat. Mech.*, 1104:P04007, 2011. arXiv:1101.2885 [math-ph].

[26] G James and M Liebeck. *Representations and Characters of Groups*. Cambridge University Press, Cambridge, 2001.

[27] T Y Lam. *A First Course in Noncommutative Rings*, volume 131 of *Graduate Texts in Mathematics*. Springer, New York, 1991.

[28] R Pierce. *Associative Algebras*, volume 88 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.

[29] M Hazewinkel, N Gubareni, and V Kirichenko. *Algebras, Rings and Modules. Vol. 1*, volume 575 of *Mathematics and its Applications*. Kluwer, New York, 2004.

[30] T Y Lam. *Lecture on Modules and Rings*, volume 189 of *Graduate Texts in Mathematics*. Springer, New York, 1998.